

Heterotopic Hand Reconstruction for Catastrophic Hand

**Santiago González Gómez¹, Emmanuel Alejandro Saucedo Resendiz², Rafael Chávez Contreras³,
María Fernanda Páez Arteaga⁴**

^{1,2}Centro Médico Nacional del Noreste, IMSS, Monterrey, Nuevo León.

¹<https://orcid.org/0009-0002-1580-5576>

²<https://orcid.org/0009-0007-9453-823X>

³Hospital Civil de Guadalajara Fray Antonio Alcalde, Guadalajara, Jalisco. <https://orcid.org/0009-0002-2298-527X>

⁴Centro Médico Nacional del Noreste, IMSS, Monterrey, Nuevo León. <https://orcid.org/0009-0005-7483-1979>

ABSTRACT

Heterotopic hand reconstruction is a surgical technique involving the transfer of amputated or non-replantable segments to alternative positions to optimize hand function following severe or mutilating injuries. This approach is employed when direct reimplantation is unfeasible due to extensive tissue damage or tissue loss at the original site. It allows for the repurposing of non-viable digits or tissues—for example, converting a non-salvageable finger into a functional thumb or utilizing parts from damaged digits to reconstruct joints, bones, or soft tissues. The choice of tissues and transfer sites is customized based on injury characteristics and functional goals, with an emphasis on restoring opposition and prehension. Clinical outcomes demonstrate that heterotopic procedures can provide significant functional gains, including sensate, power grip, and refined prehension, thus playing a crucial role in complex hand trauma reconstruction. These techniques offer effective solutions to restore hand functionality and improve patient quality of life in challenging cases.

KEYWORDS: Heterotopic hand reconstruction, finger transfer, non-replantable digit, tissue transfer, hand trauma, digital transplantation, reconstructive hand surgery, microsurgery, functional hand restoration, opposition, grip, pulley transfer, "spare parts" surgery, complex hand injury, vascularized joint transfer, reconstructive microsurgery, amputation, limb salvage, functional recovery.

ARTICLE DETAILS

Published On:
18 June 2025

Available on:
<https://ijmscrs.com/>

INTRODUCTION

Heterotopic hand reconstruction refers to the surgical transfer and replantation of amputated or non-salvageable parts of the hand (such as fingers or joints) to a different, non-anatomic location on the same or contralateral hand, with the goal of maximizing functional recovery after severe or mutilating hand injuries. This approach is typically considered when direct replantation of the amputated part to its original site is not feasible due to extensive tissue damage, contamination, or loss of the recipient site, but the amputated part or its components remain viable for transfer.

Common scenarios include:

- Transferring a non-salvageable finger or thumb to reconstruct a more functionally important digit, such as using

a finger to reconstruct a thumb, which is critical for opposition and prehension.[1-4]

- Using vascularized joints, bone, or soft tissue from amputated or non-replantable digits to reconstruct defects in other parts of the hand, thereby restoring key functions such as pinch, grip, or webspace.[2][5]
- "Spare parts" reconstruction, where tissues from non-replantable digits are recycled as grafts or flaps to address composite defects.[2][5]

The primary objective is to restore essential hand functions, particularly thumb opposition and the ability to perform tripod or lateral pinch, by prioritizing the reconstruction of a thumb and at least two opposing fingers.[1-3] Heterotopic procedures may involve microsurgical techniques, including

Heterotopic Hand Reconstruction for Catastrophic Hand

free tissue transfer and microvascular anastomosis, and can be performed acutely or in staged reconstructions.[1-2][5] Clinical outcomes reported in the literature indicate that heterotopic hand reconstruction can achieve satisfactory restoration of sensate, prehensile function, and acceptable hand appearance, with patients regaining the ability to perform activities of daily living and work-related tasks.[1-4] The choice of which digit or tissue to transfer, and to which location, is individualized based on the pattern of injury, available tissues, and functional priorities.[2-3]

Specific indications include:

- Irreparable loss or severe damage to the thumb, where a non-thumb digit (e.g., index, middle, or ring finger) is transferred to the thumb position to restore opposition and grasp.[1-2]
- Multiple digital amputations with some digits or parts non-salvageable at their original sites but suitable for transfer to more functionally advantageous positions, such as

reconstructing at least two opposable fingers in addition to the thumb.[1-2]

- Severe metacarpophalangeal or proximal interphalangeal joint destruction, where vascularized joint transfer from a non-salvageable digit is used to reconstruct a more critical ray or joint, thereby improving overall hand function.[3]
- Complex injuries where "spare parts" from non-replantable or less functional digits are recycled to reconstruct bone, skin, or web spaces, or to provide sensate coverage.[2]

In all cases, the decision to pursue heterotopic reconstruction is individualized, with priority given to restoring a functional thumb and at least two fingers capable of opposition, as this configuration is critical for meaningful hand function.[1-2] The literature emphasizes that these procedures are essential tools in the reconstructive armamentarium for maximizing functional outcomes after severe hand trauma.[3]

Table 1. Key aspects of heterotopic hand reconstruction:

Aspect	Details
Definition	Transfer of non-replantable or amputated parts to alternative hand sites
Purpose	Maximize hand function, especially opposition and prehension
Common Procedures	Digit-to-digit transfer (e.g., finger to thumb), joint or tissue transfer from non-replantable parts
Indications	Severe digital or hand injuries with non-salvageable tissues, complex trauma, or amputations
Goals	Restoring opposition, grip, pinch, range of motion, sensory function
Techniques Used	Microsurgical free tissue transfer, vascularized joint transfer, grafting
Key Tissues Transferred	Fingers, joints, soft tissues, skin from non-viable parts
Clinical Outcomes	Improved hand function, ability to perform daily activities, better aesthetic result
Advantages	Recycles tissue, restores critical functions, individualized approach
Challenges	Technical complexity, matching tissues, postoperative care requirements

Key aspects on surgical techniques

1. Heterotopic Digital Replantation: This involves replanting an amputated digit to a different position than its original anatomical site, most commonly to reconstruct a thumb or to restore opposition and pinch. For example, a non-salvageable finger may be replanted at the thumb position or another ray to maximize function. Microvascular anastomosis is required for revascularization, and the selection of recipient vessels and nerves is dictated by the defect and available tissues. Restoration of at least a thumb and two fingers is often targeted to achieve tripod pinch and basic hand function.[2, 3]
2. Heterotopic Vascularized Joint Transfer: When joints are irreparably damaged, vascularized joints from non-salvageable digits can be transferred to reconstruct critical joints (e.g., transferring a proximal interphalangeal joint from an amputated finger to a thumb or another finger). These can be performed as free tissue transfers with microvascular anastomosis or as pedicled transfers if the vascular pedicle is intact. The goal is to restore motion at key articulations, with careful attention to tendon and soft tissue reconstruction.[3]
3. Spare Parts Reconstruction: Non-viable or non-replantable tissues (bone, skin, tendon, neurovascular bundles) from amputated parts can be used as composite grafts or fillet flaps

- to reconstruct defects elsewhere in the hand. This may include using bone and skin from a non-replantable digit to reconstruct a metacarpal or webspace, or using a neurovascular fillet flap to resurface critical areas.[5]
4. Pollicization and Ray Transposition: In cases where the thumb is lost and cannot be replanted, pollicization of an index or middle finger (transferring and rotating the finger to the thumb position) may be performed, sometimes in combination with heterotopic replantation of other digits to optimize hand function.[2]
 5. Composite Tissue Flap Transfers: In complex injuries, composite tissue flaps (e.g., toe transfers, anterolateral thigh flaps, reverse posterior interosseous artery flaps) may be used to reconstruct missing digits or soft tissue, often in combination with heterotopic techniques to maximize function and coverage.[6, 7]
 6. Targeted Muscle Reinnervation and Free Flap Coverage: In select cases, targeted muscle reinnervation and free fasciocutaneous flaps may be used in conjunction with heterotopic replantation to optimize both coverage and functional outcomes, particularly in complex or bilateral injuries.[8]
- The surgical approach is dictated by the specific pattern of injury, the viability of tissues, and the reconstructive priorities

Heterotopic Hand Reconstruction for Catastrophic Hand

(e.g., restoration of sensate pinch, opposition, and grasp). The literature emphasizes the importance of individualized planning, meticulous microsurgical technique, and a staged

approach when necessary to optimize both function and aesthetics.[6-8]



Figure 1. Third finger reconstruction with inguinal flap



Figure 2. Dissection for taking the first finger for transfer, tendon dissection, dorsal artery preservation and fatty tissue preservation.



Figure 3. Successful toe to hand transfer, 1st finger left hand, tendon vein artery nerve + skin graft.

CONCLUSION

Heterotopic hand reconstruction is a valuable surgical approach in cases of severe or mutilating injuries where conventional replantation of the amputated segment is not possible. This technique allows for the transfer of non-replantable parts or tissues to alternative locations on the hand, aiming to restore critical functions such as opposition, grip, and pinch. The procedures are tailored individually based on the pattern of injury, available tissues, and the patient's functional priorities. The literature supports that these surgeries can achieve meaningful functional restoration, enabling patients to perform daily activities and work-related tasks. Overall, heterotopic reconstruction is a vital tool in the reconstructive armamentarium for maximizing hand function after complex trauma.

REFERENCES

- I. Ma Z, Zhang X, Wang G, et al. Long-Term Follow-Up of Thumb Reconstruction With a Heterotopic Replanted Finger: Case Report and Literature Review. *Frontiers in Bioengineering and Biotechnology*. 2024;12:1465108. doi:10.3389/fbioe.2024.1465108.
- II. Kokkoli E, Spyropoulou GA, Shih HS, Feng GM, Jeng SF. Heterotopic Procedures in Mutilating Hand Injuries: A Synopsis of Essential Reconstructive Tools. *Plastic and Reconstructive Surgery*. 2015;136(5):1015-1026. doi:10.1097/PRS.0000000000001721.
- III. An PC, Kuo YR, Lin TS, Yeh MC, Jeng SF. Heterotopic Replantation in Mutilating Hand Injury. *Annals of Plastic Surgery*. 2003;50(2):113-8; discusión118-9. doi:10.1097/01.SAP.0000037269.62980.69.
- IV. Tosti R, Treiser MD, Eberlin KR. Contralateral Heterotopic Thumb-to-Thumb Replantation With Free Ulnar Forearm Fasciocutaneous Flap and Targeted Muscle Reinnervation. *The Journal of Hand Surgery*. 2020;45(1):71.e1-71.e5. doi:10.1016/j.jhsa.2019.03.004.
- V. Chang NJ, Chang JT, Hsu CC, et al. Heterotopic Vascularized Joint Transfer in Mutilating Hand Injuries. *Annals of Plastic Surgery*. 2016;76 Suppl 1:S1-7. doi:10.1097/SAP.0000000000000712.
- VI. Güntürk ÖB, Kaplan İ, Yıldırım T, et al. Reconstruction of Mutilating Hand Injuries by Microsurgical Free Tissue Transfers From the Foot. *Injury*. 2021;52(12):3646-3652. doi:10.1016/j.injury.2021.03.044.
- VII. Li X, Cui J, Maharjan S, et al. Neo-Digit Functional Reconstruction of Mutilating Hand Injury Using Transplantation of Multiple Composite Tissue Flaps. *Medicine*. 2016;95(27):e4179. doi:10.1097/MD.00000000000004179.
- VIII. Tosti R, Treiser MD, Eberlin KR. Contralateral Heterotopic Thumb-to-Thumb Replantation With Free Ulnar Forearm Fasciocutaneous Flap and Targeted Muscle Reinnervation. *The Journal of Hand Surgery*. 2020;45(1):71.e1-71.e5. doi:10.1016/j.jhsa.2019.03.004.