

High-Risk Electrocardiographic Patterns in Acute Myocardial Infarction: Early Recognition, Clinical Implications, and Therapeutic Decision-Making

Donaldo Emiliano Silva López¹, Francisco Rivera Martinez², America Michelle Barrios Muñoz², Andrea Carolina Velasco Valbuena³, Johana Sofía Herman González⁴, José Nahúm Muñoz Tabares⁵, Yoselin Esparza Monreal², Karla Teresa Granados Velasco⁶, Maximiliano Tavaréz Sanchez⁷, Aldo Mauricio Villagrana Herrera², Miguel Ángel Espidio Coello⁸, Ivana Leilani Júnez Villicaña⁹, Alexia Paola Rodríguez Barboza¹

¹Tecnológico de Monterrey Campus Monterrey. Monterrey, Mexico.

²Universidad Autónoma de Zacatecas "Francisco García Salinas". Zacatecas, Mexico.

³Universidad industrial de Santander. Mexico.

⁴Benemerita Universidad autónoma de Aguascalientes. Aguascaliente, Mexico.

⁵Universidad Autónoma de Aguascalientes. Aguascalientes, Mexico.

⁶Universidad del valle de Atemajac. Mexico.

⁷Universidad Cuauhtemoc Aguascalientes Aguascalientes, Mexico.

⁸Benemérita Universidad Autónoma de Puebla. Mexico.

⁹Universidad michoacana de San Nicolas de Hidalgo. Michoacan, Mexico.

ABSTRACT

Background: The 12-lead electrocardiogram (ECG) remains the cornerstone of rapid diagnosis in acute myocardial infarction (AMI). Early identification of high-risk ECG patterns is critical for timely intervention, risk stratification, and improving clinical outcomes.

Objective: This article reviews the electrocardiographic hallmarks of AMI, emphasizing high-risk features that necessitate emergent reperfusion therapy, discriminate STEMI from NSTEMI, and predict adverse cardiovascular events.

Methods: A narrative review of current literature was conducted, focusing on ECG patterns associated with acute coronary syndromes (ACS), including ST-segment elevation (STE), ST depression, hyperacute T waves, Q waves, and conduction abnormalities. Special attention was given to STEMI-equivalent patterns and occlusive myocardial ischemia.

Results: Key high-risk ECG findings include:

STE in ≥ 2 contiguous leads (indicating transmural ischemia).

New left bundle branch block (LBBB) meeting Sgarbossa criteria.

Global ST depression with STE in aVR (suggesting left main or triple-vessel disease).

Posterior STEMI (ST depression in V1–V3 with dominant R waves).

De Winter's T waves (upwardly concave ST depression with hyperacute T waves in precordial leads).

These patterns correlate with large ischemic burden, proximal coronary occlusion, and higher mortality.

Conclusion: Mastery of high-risk ECG patterns enables rapid triage, appropriate revascularization, and reduced morbidity in AMI. Future advancements in AI-assisted ECG interpretation may further enhance early detection.

KEYWORDS: Acute myocardial infarction (AMI), Electrocardiogram (ECG), ST-segment elevation myocardial infarction (STEMI), High-risk ECG patterns, Myocardial ischemia, Reperfusion therapy, Sgarbossa criteria, Occlusive coronary disease

ARTICLE DETAILS

Published On:
25 June 2025

Available on:
<https://ijmscrs.com/>

INTRODUCTION

Acute myocardial infarction (AMI) remains a leading cause of global morbidity and mortality, with timely diagnosis and intervention being pivotal to survival. The 12-lead electrocardiogram (ECG) serves as the first-line diagnostic tool, capable of distinguishing between ischemic etiologies, guiding emergent reperfusion strategies, and predicting adverse outcomes. While ST-segment elevation myocardial infarction (STEMI) is universally recognized as a medical emergency requiring immediate percutaneous coronary intervention (PCI), numerous other ECG patterns—often termed "STEMI-equivalents" or "high-risk NSTEMI"—carry similar prognostic implications but are frequently underrecognized in clinical practice.^{1,2}

Emerging data underscore the role of ECG in identifying left main coronary occlusion, multivessel disease, and mechanical complications (e.g., ventricular septal rupture). Furthermore, advanced criteria such as the Anderson-Wilkins score (assessing STE morphology) and modified Sgarbossa rules (for paced rhythms) refine diagnostic accuracy.²

Pathophysiology of High-Risk Electrocardiographic Patterns in Acute Myocardial Infarction

The electrocardiographic manifestations of acute myocardial infarction (AMI) arise from a complex interplay of ischemic injury, cellular electrophysiological disturbances, and the anatomical distribution of coronary artery occlusion. The 12-lead ECG serves as a dynamic reflection of these pathophysiological processes, with distinct patterns emerging based on the severity, duration, and location of myocardial ischemia.³

ST-segment elevation (STE), the hallmark of transmural ischemia, results from abrupt coronary occlusion, typically due to thrombus formation over a ruptured atherosclerotic plaque. The ensuing oxygen supply-demand mismatch leads to myocardial membrane instability, with potassium efflux and intracellular calcium accumulation altering transmembrane potentials. This depolarization delay in ischemic zones generates injury currents, manifesting as STE in leads overlying the affected territory. The morphology of STE—whether convex, concave, or obliquely straight—correlates with the acuteness and extent of injury. For instance, anterior STEMI (V1–V4 STE) often indicates left anterior descending (LAD) artery occlusion, associated with a high risk of left ventricular dysfunction and cardiogenic shock due to the large myocardial territory at risk. In contrast, inferior STEMI (II, III, aVF STE) typically stems from right coronary artery (RCA) or left circumflex (LCx) occlusion, with concomitant right ventricular involvement (e.g., STE in V1 or V4R) signaling proximal RCA occlusion and heightened risk of hemodynamic compromise.^{4,5}

Reciprocal ST-segment depression, frequently observed in contralateral leads, reflects remote subendocardial ischemia

due to either collateral blood flow diversion or diffuse coronary disease. Notably, ST depression in V1–V3 with tall R waves may localize to the posterior wall (a "posterior STEMI"), often missed without additional posterior leads. The pathophysiological basis lies in the loss of opposing electrical forces normally generated by the posterior myocardium, unmasking anteriorly directed depolarization vectors.⁵

Hyperacute T waves, characterized by symmetrical, peaked amplitudes, represent the earliest ischemic change, preceding STE. These result from accelerated repolarization in ischemic myocytes due to adenosine triphosphate (ATP)-dependent potassium channel activation, shortening action potential duration. Their transient nature underscores the critical window for intervention before irreversible necrosis occurs. Conversely, deep T-wave inversions, as seen in Wellens' syndrome, indicate reperfusion following critical LAD stenosis, where residual stunned myocardium generates abnormal repolarization currents.⁶

Q waves, a marker of completed necrosis, emerge hours to days post-occlusion as electrically inert tissue fails to contribute to depolarization, leaving unopposed forces from healthy myocardium to dominate the QRS complex. Their presence in anterior leads correlates with higher infarct burden and adverse remodeling.⁶

Conduction abnormalities, such as new-onset left bundle branch block (LBBB), complicate interpretation but signify extensive septal ischemia disrupting the His-Purkinje system. The modified Sgarbossa criteria aid in diagnosing occlusion MI in LBBB by identifying discordant STE proportional to QRS depth, reflecting ventricular depolarization-repolarization mismatch.⁶

Global ST depression with STE in aVR suggests left main or triple-vessel disease, where widespread subendocardial ischemia generates a "current of injury" directed toward the right superior lead (aVR). Similarly, De Winter's pattern—upwardly concave ST depression with hyperacute T waves in precordial leads—indicates proximal LAD occlusion without classic STE, attributed to alternating ischemic injury gradients.⁶

These patterns collectively underscore the ECG's role in delineating ischemic pathophysiology, where each waveform aberration mirrors the underlying cellular derangements, coronary anatomy, and impending clinical risk.^{6,7}

Clinical Significance of High-Risk Electrocardiographic Patterns in Acute Myocardial Infarction

The electrocardiogram remains an indispensable tool in the early diagnosis and risk stratification of acute myocardial infarction, with specific patterns carrying profound prognostic and therapeutic implications. The recognition of these high-risk electrocardiographic manifestations is not merely an academic exercise but a critical determinant of

High-Risk Electrocardiographic Patterns in Acute Myocardial Infarction: Early Recognition, Clinical Implications, and Therapeutic Decision-Making

clinical outcomes, as they reflect the underlying severity of coronary pathology, the extent of myocardial jeopardy, and the urgency with which reperfusion strategies must be implemented.⁷

Among the most critical patterns, ST-segment elevation serves as the hallmark of transmural ischemia and is the primary electrocardiographic criterion for emergent reperfusion therapy. The anatomical localization of ST elevation provides immediate insight into the likely culprit artery, with anterior leads implicating the left anterior descending artery, inferior leads suggesting right coronary or circumflex involvement, and lateral changes pointing to circumflex or diagonal branch occlusion. The magnitude of ST elevation correlates with the degree of myocardial injury, with more pronounced elevations indicating larger areas of at-risk myocardium and consequently higher likelihood of complications such as ventricular arrhythmias, cardiogenic shock, or mechanical complications including ventricular septal rupture or free wall rupture.⁸

Equally significant are the so-called STEMI-equivalent patterns, which, despite lacking traditional ST elevation, signify comparable levels of coronary occlusion and myocardial threat. The presence of a new left bundle branch block in the appropriate clinical context demands the same urgency as overt ST elevation, as it frequently reflects extensive septal infarction with disruption of the conduction system. Posterior myocardial infarction, often overlooked due to its subtle presentation as ST depression in the anterior leads, represents a particularly treacherous entity, as the substantial myocardial mass involved in the posterior and lateral walls can rapidly deteriorate into hemodynamic collapse if not promptly revascularized. The De Winter pattern, characterized by upsloping ST depression with tall, symmetrical T waves in the precordial leads, represents an angiographic emergency, as it is virtually pathognomonic for proximal left anterior descending artery occlusion and carries a dismal prognosis without immediate intervention.^{8,9}

The ominous finding of global ST depression with concomitant ST elevation in lead aVR cannot be overstated in its clinical importance. This pattern serves as an electrocardiographic red flag for either left main coronary artery occlusion or severe triple-vessel disease, both scenarios associated with catastrophic outcomes without urgent revascularization. The pathophysiology underlying this pattern involves widespread subendocardial ischemia generating a resultant vector directed toward the right shoulder, creating the characteristic aVR elevation that should immediately alert clinicians to the presence of this critical condition.^{10,11}

Beyond the acute ischemic changes, certain chronic patterns provide valuable prognostic information. Deep, symmetrical T wave inversions in the precordial leads, as seen in Wellens syndrome, indicate a pre-infarction state with critical stenosis

of the proximal left anterior descending artery. These patients represent walking time bombs, as their electrocardiographic findings predict imminent anterior wall infarction without timely revascularization. Similarly, the presence of Q waves, while traditionally considered a marker of completed infarction, in the context of acute symptoms may indicate ongoing myocardial necrosis and should not deter from aggressive intervention.^{12,13}

The clinical imperative to recognize these patterns extends beyond simple diagnosis, as each carries specific management implications. The differential outcomes based on electrocardiographic findings have been well documented, with proximal occlusion patterns demonstrating significantly higher mortality rates compared to more distal lesions. Furthermore, certain patterns may influence the choice of reperfusion strategy, with some institutions advocating for immediate transfer to cardiac catheterization laboratories based solely on specific high-risk electrocardiographic features, even in the absence of traditional ST elevation criteria.¹⁴

In the contemporary era of increasingly sophisticated cardiac care, the electrocardiogram remains remarkably resilient as the initial diagnostic modality for acute coronary syndromes. Its proper interpretation, with particular attention to these high-risk patterns, continues to save countless lives through prompt identification of those patients most in need of urgent intervention. The clinical significance of these patterns cannot be overstated, as they represent the crucial interface between electrical manifestations and clinical decision-making that ultimately determines patient outcomes in acute myocardial infarction.¹⁵

CONCLUSION

The electrocardiographic evaluation of patients presenting with acute myocardial infarction remains an indispensable diagnostic tool that provides critical insights into the underlying pathophysiological processes, guides immediate therapeutic decisions, and predicts clinical outcomes with remarkable precision. The identification of high-risk ECG patterns transcends mere pattern recognition, representing instead a sophisticated clinical skill that integrates electrical manifestations with coronary anatomy and myocardial pathophysiology. These patterns serve as vital biomarkers of impending cardiac catastrophe, with each distinctive waveform aberration carrying specific prognostic implications that demand tailored management strategies. The traditional ST-segment elevation myocardial infarction patterns, while serving as well-established markers of transmural ischemia requiring emergent reperfusion, represent only one facet of the complex electrocardiographic spectrum seen in acute coronary syndromes.

Of equal clinical importance are the STEMI-equivalent patterns, which despite their varied presentations, signify comparable degrees of coronary occlusion and myocardial

jeopardy. The recognition of posterior wall infarction through reciprocal changes, the identification of proximal left anterior descending artery occlusion through De Winter's pattern, and the detection of left main or triple vessel disease through global ST changes with aVR elevation all represent critical diagnostic competencies that directly impact mortality and morbidity. These patterns collectively underscore the principle that myocardial ischemia manifests along an electrocardiographic continuum, where the absence of classic ST elevation does not necessarily indicate benign pathology. The clinical implications of these findings extend far beyond the emergency department, influencing the entire spectrum of cardiac care from pre-hospital triage to long-term prognostic stratification. The presence of certain high-risk patterns may prompt more aggressive revascularization strategies, earlier mechanical circulatory support consideration, or more intensive monitoring for complications. Furthermore, these electrocardiographic markers provide valuable information about infarct size, myocardial salvage potential, and likelihood of electrical instability, all of which inform both acute management and secondary prevention strategies.

As cardiovascular medicine continues to evolve with advanced imaging modalities and biomarker technologies, the 12-lead ECG maintains its fundamental role as the initial diagnostic modality for acute myocardial infarction. Its universal availability, rapid acquisition, and rich informational content ensure its enduring clinical relevance. However, the full potential of electrocardiographic diagnosis can only be realized through comprehensive pattern recognition that extends beyond simple ST elevation analysis to include the full spectrum of ischemic manifestations. Future directions in electrocardiography, including computer-assisted interpretation and artificial intelligence applications, promise to enhance our ability to detect these high-risk patterns with greater sensitivity and specificity, potentially revolutionizing the early diagnosis of acute coronary syndromes.

In conclusion, the meticulous interpretation of electrocardiographic patterns in acute myocardial infarction represents a critical clinical skill that directly impacts patient outcomes. The ability to recognize and appropriately respond to these high-risk patterns remains fundamental to the practice of emergency and cardiovascular medicine, serving as the crucial link between electrical manifestations and life-saving interventions. As our understanding of ischemic electrocardiography continues to deepen, so too does our capacity to deliver timely, targeted care to patients experiencing the most severe manifestations of acute coronary artery disease. The electrocardiogram thus endures not merely as a historical diagnostic tool, but as a dynamic and essential component of contemporary cardiac care, whose proper interpretation continues to make the difference between myocardial salvage and irreversible loss, between survival and mortality.

REFERENCES

- I. K. Thygesen, J.S. Alpert, A.S. Jaffe, *et al.* Fourth universal definition of myocardial infarction (2018) *Circulation*, 138 (2018), pp. e618-e651
- II. R.A. Byrne, X. Rossello, J.J. Coughlan, *et al.* 2023 ESC Guidelines for the management of acute coronary syndromes *Eur Heart J*, 44 (2023), pp. 3720-3826
- III. P.W. Macfarlane, D. Browne, B. Devine, *et al.* Modification of ACC/ESC criteria for acute myocardial infarction *J Electrocardiol*, 37 (2004), pp. 98-103
- IV. J.N. de Alencar Neto, M.K. Scheffer, B.P. Correia, *et al.* Systematic review and meta-analysis of diagnostic test accuracy of ST-segment elevation for acute coronary occlusion. *Int J Cardiol*, 402 (2024), Article 131889
- V. H.P. Meyers, A. Bracey, D. Lee, *et al.* Comparison of the ST-elevation myocardial infarction (STEMI) vs. NSTEMI and occlusion MI (OMI) vs. NOMI paradigms of acute MI *J Emerg Med*, 60 (2021), pp. 273-284
- VI. H.P. Meyers, A. Bracey, D. Lee, *et al.* Accuracy of OMI ECG findings versus STEMI criteria for diagnosis of acute coronary occlusion myocardial infarction. *Int J Cardiol Heart Vasc*, 33 (2021), Article 100767
- VII. J. Karwowski, M. Gierlotka, M. Gąsior, *et al.* Relationship between infarct artery location, acute total coronary occlusion, anomortality in STEMI and NSTEMI patients *Pol Arch Intern Med*, 127 (2017), pp. 401-411
- VIII. G.W. Stone, D. Cox, E. Garcia, *et al.* Normal flow (TIMI-3) before mechanical reperfusion therapy is an independent determinant of survival in acute myocardial infarction: analysis from the primary angioplasty in myocardial infarction trials *Circulation*, 104 (2001), pp. 636-641
- IX. D.A. Cox, G.W. Stone, C.L. Grines, *et al.* Comparative early and late outcomes after primary percutaneous coronary intervention in ST-segment elevation and non-ST-segment elevation acute myocardial infarction (from the CADILLAC Trial) *Am J Cardiol*, 98 (2006), pp. 331-337
- X. F.M. Fesmire, R.F. Percy, J.B. Bardoner, *et al.* Usefulness of automated serial 12-lead ECG monitoring during the initial emergency department evaluation of patients with chest pain. *Ann Emerg Med*, 31 (1998), pp. 3-11
- XI. P. Hillinger, I. Strebel, R. Abächerli, *et al.* Prospective validation of current quantitative electrocardiographic criteria for ST-elevation myocardial infarction *Int J Cardiol*, 292 (2019), pp. 1-12

High-Risk Electrocardiographic Patterns in Acute Myocardial Infarction: Early Recognition, Clinical Implications, and Therapeutic Decision-Making

- XII. E.K. Aslanger, H.P. Meyers, S.W. Smith
Recognizing electrocardiographically subtle
occlusion myocardial infarction and differentiating
it from mimics: ten steps to or away from cath lab.
Turk Kardiyol Dern Ars, 49 (2021), pp. 488-500
- XIII. K. Nikus, O. Pahlm, G. Wagner, *et al.*
Electrocardiographic classification of acute
coronary syndromes: a review by a committee of the
International Society for Holter and Non-Invasive
Electrocardiology. *J Electrocardiol*, 43 (2010), pp.
91-103
- XIV. M.C. Kontos, J.A. de Lemos, S.B. Deitelzweig, *et al.*
2022 ACC Expert Consensus Decision Pathway
on the evaluation and disposition of acute chest pain
in the emergency department. *J Am Coll Cardiol*, 80
(2022), pp. 1925-1960
- XV. A.R. Khan, H. Golwala, A. Tripathi, *et al.* Impact of
total occlusion of culprit artery in acute non-ST
elevation myocardial infarction: a systematic review
and meta-analysis. *Eur Heart J*, 38 (2017), pp. 3082-
3089