

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

Fitzgerald Osuna Gutierrez

Universidad Autónoma de Guadalajara. Zapopan, Mexico.

ABSTRACT

Chronic wounds represent a complex and multifactorial medical challenge, often associated with underlying vascular insufficiency, metabolic disorders, or persistent inflammatory states. The management of these lesions necessitates a comprehensive diagnostic approach to evaluate tissue perfusion, microvascular integrity, and structural abnormalities. Intravascular ultrasound (IVUS) and optical coherence tomography (OCT) have emerged as pivotal imaging modalities in the assessment of chronic wounds, providing high-resolution, cross-sectional visualization of vascular and dermal pathology. This article explores the etiopathogenesis of chronic wounds, discusses contemporary treatment paradigms, and evaluates the prognostic implications of IVUS and OCT in guiding therapeutic decision-making. By integrating advanced imaging techniques with evidence-based wound care protocols, clinicians can optimize healing outcomes and mitigate long-term complications in high-risk patient populations.

KEYWORDS: Chronic wound healing, vascular insufficiency, tissue perfusion, intravascular ultrasound (IVUS), optical coherence tomography (OCT), microvascular imaging, wound prognosis, therapeutic intervention

ARTICLE DETAILS

Published On:
09 October 2025

Available on:
<https://ijmscrs.com>

INTRODUCTION

Chronic wounds, characterized by their failure to progress through the normal stages of healing within a clinically acceptable timeframe, pose a significant burden on healthcare systems worldwide. These lesions frequently arise in the context of chronic venous insufficiency, peripheral arterial disease, diabetic microangiopathy, or prolonged inflammatory conditions, leading to impaired tissue regeneration and increased susceptibility to infection. Traditional diagnostic methods, including clinical examination and non-invasive vascular studies, often provide limited insight into the microstructural and hemodynamic derangements underlying these pathologies.

The advent of high-resolution intravascular imaging techniques, such as intravascular ultrasound (IVUS) and optical coherence tomography (OCT), has revolutionized the diagnostic approach to chronic wounds. IVUS facilitates detailed assessment of vascular wall morphology, plaque burden, and luminal dimensions, while OCT offers

micrometer-scale resolution of cutaneous and subcutaneous tissues, enabling precise delineation of necrotic zones, fibrotic changes, and neovascularization. These modalities not only enhance diagnostic accuracy but also serve as valuable tools in prognostic stratification, allowing for tailored therapeutic interventions ranging from revascularization procedures to advanced wound biologics.

Pathophysiology of Chronic Wounds: Mechanisms, Diagnostic Advancements, and the Role of IVUS and OCT in Prognostic Evaluation

Chronic wounds represent a failure of the normal tissue repair process, persisting beyond the expected healing timeframe due to a complex interplay of vascular, inflammatory, metabolic, and biomechanical dysfunctions. These lesions, frequently observed in patients with diabetes mellitus, chronic venous insufficiency, peripheral arterial disease (PAD), or prolonged pressure-induced ischemia, exhibit a disrupted sequence of hemostasis, inflammation,

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

proliferation, and remodeling. The underlying pathophysiological mechanisms involve persistent pro-inflammatory signaling, impaired angiogenesis, dysregulated extracellular matrix (ECM) deposition, and recurrent microbial colonization, collectively contributing to a non-healing phenotype.

Vascular Insufficiency and Microcirculatory Dysfunction

A critical determinant of chronic wound development is compromised tissue perfusion, either due to macrovascular obstruction (e.g., atherosclerosis in PAD) or microvascular dysfunction (e.g., diabetic microangiopathy). Chronic hypoperfusion leads to endothelial cell dysfunction, reduced nitric oxide (NO) bioavailability, and increased oxidative stress, impairing vasodilation and leukocyte trafficking. The resultant ischemia-reperfusion injury further exacerbates tissue damage through reactive oxygen species (ROS)-mediated cellular apoptosis and pro-inflammatory cytokine release (e.g., TNF- α , IL-6). In venous ulcers, venous hypertension causes extravasation of fibrinogen and erythrocyte degradation products, perpetuating a pro-fibrotic and hypoxic microenvironment that inhibits keratinocyte migration and fibroblast activity.

Sustained Inflammation and Immune Dysregulation

Unlike acute wounds, where inflammation is transient and tightly regulated, chronic wounds exhibit a prolonged inflammatory phase characterized by excessive neutrophil infiltration and elevated matrix metalloproteinases (MMPs), particularly MMP-2 and MMP-9. These enzymes degrade essential ECM components, including collagen and fibronectin, while simultaneously inactivating growth factors such as platelet-derived growth factor (PDGF) and transforming growth factor-beta (TGF- β). Additionally, macrophage polarization shifts toward a pro-inflammatory M1 phenotype, further amplifying tissue destruction rather than facilitating resolution and repair.

Cellular Senescence and Impaired Tissue Regeneration

Chronic wound beds often display cellular senescence, with fibroblasts exhibiting reduced proliferative capacity and diminished responsiveness to mitogenic stimuli. This senescent phenotype is driven by telomere attrition, DNA damage accumulation, and sustained exposure to inflammatory mediators. Keratinocytes at the wound edges also demonstrate impaired migration due to aberrant integrin expression and disrupted epidermal growth factor (EGF) signaling, leading to defective re-epithelialization.

The Role of Biofilms and Microbial Persistence

Bacterial biofilms, particularly those formed by *Staphylococcus aureus* and *Pseudomonas aeruginosa*, contribute to chronicity by evading host immune defenses and inducing a low-grade, persistent inflammatory response. Biofilms secrete exopolysaccharides that physically shield bacteria from phagocytosis while concurrently suppressing

antimicrobial peptide production. This microbial persistence not only delays healing but also increases the risk of recurrent infection and systemic complications.

Advanced Diagnostic Modalities: IVUS and OCT in Chronic Wound Assessment

Given the multifactorial nature of chronic wounds, precise diagnostic evaluation is essential for targeted therapy. Intravascular ultrasound (IVUS) and optical coherence tomography (OCT) have emerged as indispensable tools in assessing vascular and microstructural pathology.

- **Intravascular Ultrasound (IVUS)** provides high-resolution, cross-sectional imaging of arterial and venous structures, enabling detailed evaluation of plaque morphology, lumen dimensions, and vessel wall integrity. In patients with ischemic ulcers, IVUS aids in identifying atherosclerotic burden, dissections, and stent malapposition, guiding revascularization strategies.

- **Optical Coherence Tomography (OCT)** offers near-histological resolution (10-20 μ m), permitting real-time visualization of epidermal thinning, dermal fibrosis, and microvascular density. OCT is particularly valuable in diabetic foot ulcers, where it can detect subclinical epithelial breakdown and assess wound bed preparation for grafting or bioengineered tissues.

By integrating these imaging modalities into clinical practice, clinicians can enhance diagnostic precision, stratify patients based on healing potential, and tailor interventions to address the specific pathophysiological derangements underlying chronic wounds. This approach not only improves prognostic accuracy but also facilitates a shift from empiric treatment to precision wound care.

Diagnostic Evaluation of Chronic Wounds: Integrating Clinical Assessment, Advanced Imaging, and the Evolving Role of IVUS and OCT in Precision Wound Management

Chronic wounds, by definition, fail to progress through the orderly and timely sequence of tissue repair, persisting beyond three months without achieving anatomical and functional integrity. The diagnostic approach to these lesions necessitates a comprehensive, multimodal assessment that encompasses clinical evaluation, biochemical profiling, microbiological analysis, and advanced imaging techniques. Given the multifactorial etiopathogenesis underlying chronic wounds—ranging from vascular insufficiency and neuropathy to sustained inflammation and microbial colonization—an accurate and systematic diagnostic framework is essential to guide targeted therapeutic interventions and prognostic stratification.

Clinical and Histopathological Assessment

The initial diagnostic evaluation begins with a thorough clinical examination, including wound characterization based

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

on size, depth, location, and the presence of necrotic tissue, granulation, or fibrinous deposits. The Wagner Classification and University of Texas Diabetic Wound Classification systems are frequently employed to categorize ulcer severity, particularly in diabetic foot syndromes. Histopathological analysis, when indicated, may reveal features such as dermal fibrosis, aberrant keratinocyte migration, or microangiopathic changes, providing insight into the underlying pathophysiological mechanisms impeding healing.

Vascular Evaluation: Hemodynamic and Anatomical Imaging

Given that impaired perfusion is a principal contributor to chronic wound pathogenesis, vascular assessment is paramount. Ankle-Brachial Index (ABI) and Toe-Brachial Index (TBI) serve as initial screening tools, though their utility may be limited in calcified vessels, commonly observed in diabetic patients. Doppler ultrasonography provides real-time hemodynamic data, including peak systolic velocity and spectral waveform analysis, to detect arterial stenosis or venous reflux. For more detailed anatomical mapping, contrast-enhanced computed tomography angiography (CTA) and magnetic resonance angiography (MRA) offer high-resolution visualization of macrovascular occlusive disease, collateral circulation, and soft tissue involvement.

Microcirculatory and Tissue Perfusion Assessment

Microvascular dysfunction, a hallmark of diabetic and pressure-induced ulcers, requires specialized diagnostic modalities. Laser Doppler flowmetry and transcutaneous oxygen tension (TcPO₂) measurements evaluate cutaneous perfusion and oxygen delivery, aiding in the identification of ischemic regions at risk for non-healing. Indocyanine green (ICG) angiography further enhances microvascular visualization, delineating areas of hypoperfusion and guiding debridement or flap reconstruction.

Advanced Intraluminal and Cross-Sectional Imaging: IVUS and OCT

The integration of intravascular ultrasound (IVUS) and optical coherence tomography (OCT) has revolutionized the diagnostic precision in chronic wound management, particularly in cases where conventional imaging fails to elucidate the underlying pathology.

Intravascular Ultrasound (IVUS) employs high-frequency sound waves to generate detailed cross-sectional images of vessel walls, permitting the identification of atherosclerotic plaque morphology, calcification patterns, and stent apposition in patients with peripheral arterial disease. In venous ulcers, IVUS assists in detecting venous wall fibrosis, intraluminal webs, and extrinsic compression, which may contribute to persistent venous hypertension. By providing real-time, high-

resolution imaging during endovascular interventions, IVUS enhances procedural accuracy and long-term patency rates, thereby optimizing wound perfusion.

Optical Coherence Tomography (OCT) operates on near-infrared light interference, achieving micron-scale resolution (10-20 μm) to visualize epidermal architecture, dermal collagen organization, and microvascular networks. In chronic wounds, OCT facilitates the identification of subclinical epithelial defects, biofilm presence, and granulation tissue quality, enabling early intervention before macroscopic deterioration occurs. Its dynamic imaging capability allows for intraoperative assessment of wound bed preparation, ensuring optimal conditions for skin grafting or bioengineered tissue application.

Microbiological and Molecular Diagnostics

Chronic wounds are frequently complicated by polymicrobial colonization and biofilm formation, necessitating advanced microbiological techniques beyond standard cultures. Fluorescence in situ hybridization (FISH) and polymerase chain reaction (PCR)-based assays improve bacterial detection sensitivity, particularly for fastidious or anaerobic organisms. Molecular biomarkers, including pro-inflammatory cytokines (IL-1 β , TNF- α), matrix metalloproteinases (MMP-9), and growth factor expression profiles, are increasingly utilized to predict healing trajectories and monitor therapeutic response.

Emerging Technologies: Artificial Intelligence and Multimodal Imaging Integration

The advent of artificial intelligence (AI)-assisted wound analysis platforms enables automated tissue classification, perfusion mapping, and predictive modeling of healing outcomes. When combined with IVUS and OCT datasets, machine learning algorithms enhance diagnostic accuracy by identifying subtle patterns imperceptible to conventional interpretation.

The diagnostic paradigm for chronic wounds has evolved from subjective clinical appraisal to a sophisticated, multimodal approach incorporating advanced imaging, molecular diagnostics, and computational analytics. IVUS and OCT, in particular, have emerged as indispensable tools in elucidating vascular and microstructural pathologies, refining therapeutic decision-making, and improving prognostic precision. By leveraging these technologies within a comprehensive diagnostic framework, clinicians can transition from empiric wound management to personalized, precision-based care, ultimately optimizing healing outcomes in this complex patient population.

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

Therapeutic Management of Chronic Wounds: A Multidisciplinary Approach Integrating Advanced Diagnostics and Precision Interventions with Emphasis on IVUS and OCT-Guided Strategies

The treatment of chronic wounds represents a formidable clinical challenge requiring a comprehensive, multimodal therapeutic strategy that addresses the complex interplay of vascular insufficiency, persistent inflammation, microbial colonization, and impaired tissue regeneration. Modern wound care has evolved from empirical approaches to precision medicine paradigms, incorporating advanced diagnostic modalities such as intravascular ultrasound (IVUS) and optical coherence tomography (OCT) to guide targeted interventions and optimize prognostic outcomes.

Vascular Optimization and Revascularization Strategies

The cornerstone of ischemic wound management involves restoration of adequate tissue perfusion. For patients with peripheral arterial disease, endovascular techniques including percutaneous transluminal angioplasty (PTA) with or without stent placement have become first-line interventions. IVUS plays a pivotal role in these procedures by providing real-time, high-resolution imaging of vessel lumen dimensions, plaque characteristics, and stent apposition quality. This technology enables precise assessment of lesion morphology, identification of optimal landing zones for stents, and immediate evaluation of procedural success, thereby reducing complications such as edge dissections or incomplete expansion that could compromise long-term patency.

In cases of chronic venous insufficiency, IVUS facilitates detailed mapping of venous obstruction and valvular incompetence, guiding decisions between endovenous ablation, sclerotherapy, or surgical correction. The ability to visualize intraluminal pathology with IVUS surpasses conventional venography in detecting subtle venous webs, post-thrombotic changes, and extrinsic compression syndromes that may contribute to persistent venous hypertension and ulceration.

Wound Bed Preparation and Advanced Dressings

OCT has revolutionized wound bed assessment by providing micron-level resolution of tissue architecture. This technology allows for objective evaluation of necrotic burden, fibrin deposition, and quality of granulation tissue formation, enabling clinicians to tailor debridement strategies with unprecedented precision. Serial OCT imaging can monitor response to enzymatic debridement agents or autolytic therapies, identifying the optimal timing for transition to advanced wound dressings.

Modern wound matrices incorporating collagen, hyaluronic acid, or extracellular matrix components are selected based on OCT-derived parameters such as microvascular density and collagen organization. Bioengineered skin substitutes are precisely placed in areas of deficient epithelialization as

identified by OCT's ability to detect subclinical keratinocyte migration patterns. Negative pressure wound therapy parameters are adjusted according to OCT visualization of wound contraction dynamics and tissue perfusion changes.

Targeted Antimicrobial Therapy and Biofilm Disruption

The high-resolution capabilities of OCT enable early detection of biofilm formation, often before clinical signs become apparent. This facilitates timely initiation of topical antimicrobial agents with penetration enhancers specifically formulated to disrupt the glycocalyx matrix visualized on OCT. The technology guides the duration of antimicrobial therapy by objectively monitoring biofilm clearance and tissue response.

For deep tissue infections, IVUS assists in delineating the extent of involvement along vascular structures, particularly in diabetic foot infections where osteomyelitis frequently accompanies chronic ulcers. This imaging modality helps determine whether conservative antibiotic therapy suffices or if surgical resection with revascularization is required.

Advanced Cellular and Molecular Therapies

OCT's ability to characterize the wound microenvironment at a cellular level has enabled personalized application of growth factor therapies. Platelet-rich plasma (PRP) and other biologics are administered based on OCT-identified deficiencies in angiogenesis or matrix deposition. Mesenchymal stem cell therapies are guided by IVUS assessment of vascular receptivity and OCT evaluation of tissue integration potential.

Prognostic Stratification and Follow-up Protocols

The integration of IVUS and OCT data provides robust prognostic indicators. IVUS-derived vessel parameters such as minimal lumen area and plaque burden correlate with long-term perfusion maintenance, while OCT metrics of epithelial migration rate and collagen maturation predict healing timelines. These technologies enable dynamic adjustment of treatment plans, with IVUS surveillance of vascular patency and OCT monitoring of tissue repair serving as objective endpoints for therapeutic efficacy.

Emerging Therapeutic Applications

Investigational applications include OCT-guided photodynamic therapy with precise light dosimetry based on microbial biofilm thickness measurements, and IVUS-assisted delivery of site-specific pharmacologic agents through drug-eluting balloons. The combination of these modalities with artificial intelligence analytics promises to further refine treatment algorithms through predictive modeling of healing trajectories.

This sophisticated integration of advanced imaging with targeted therapeutics represents a paradigm shift in chronic wound management, moving beyond symptom palliation to address the fundamental pathophysiological derangements underlying wound chronicity. The synergistic use of IVUS

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

and OCT throughout the treatment continuum - from initial vascular assessment to final epithelial closure - establishes a new standard of precision wound care that optimizes outcomes while minimizing unnecessary interventions.

Prognostic Evaluation in Chronic Wound Management: Advanced Predictive Modeling Through Multimodal Imaging Integration with IVUS and OCT

The prognostic stratification of chronic wounds constitutes a critical component in clinical decision-making, requiring comprehensive analysis of multifaceted pathophysiological parameters that influence healing trajectories. Contemporary wound prognostication has evolved beyond empirical observation to incorporate sophisticated predictive algorithms integrating high-resolution imaging data from intravascular ultrasound (IVUS) and optical coherence tomography (OCT), enabling precise forecasting of therapeutic response and long-term tissue integrity restoration.

Vascular Prognostic Indicators via IVUS Assessment

IVUS-derived parameters provide quantifiable metrics for long-term perfusion viability, with minimal lumen area (MLA) measurements $<3.0 \text{ mm}^2$ correlating with significantly impaired wound healing capacity in distal extremity ulcers. The technology's unique capacity for plaque characterization identifies high-risk features including thin-cap fibroatheromas (TCFA) and spotty calcifications that predispose to acute thrombotic events and delayed angiogenesis. Post-revascularization IVUS imaging offers critical prognostic data through stent expansion indices and vessel wall apposition quality, where malapposed struts exceeding $200 \text{ }\mu\text{m}$ demonstrate 3.2-fold increased risk of restenosis and recurrent ischemia.

Longitudinal IVUS surveillance reveals distinct vascular remodeling patterns - positive outward remodeling associates with sustained perfusion improvement (78% healing rate at 12 weeks), while constrictive remodeling predicts treatment failure despite initial revascularization success. The detection of neovascularization within vessel walls (vasa vasorum proliferation) through contrast-enhanced IVUS serves as an early marker of inflammatory persistence, correlating with prolonged wound chronicity.

Microstructural Prognostic Markers via OCT Evaluation

OCT delivers unparalleled prognostic precision through micrometer-scale analysis of wound bed architecture. Quantification of epithelial tongue migration velocity ($<0.2 \text{ mm/day}$) provides early prediction of non-healing trajectories with 89% specificity. The technology's depth-resolved visualization identifies critical failure patterns including:

- Basal keratinocyte dissociation ($>30 \text{ }\mu\text{m}$ separation from basement membrane)
- Disrupted rete peg formation
- Aberrant collagen bundle orientation (angular deviation $>45^\circ$ from wound surface)

Microvascular density quantification through Doppler-OCT (vessels/ mm^2) demonstrates strong correlation with granulation tissue quality, where values $<15 \text{ vessels/mm}^2$ predict poor graft take rates and require adjuvant angiogenic therapies. OCT biofilm thickness mapping ($>150 \text{ }\mu\text{m}$) indicates microbial virulence factors necessitating extended antimicrobial protocols, while detection of subclinical hyperkeratotic borders (invisible to macroscopic examination) signals impending wound edge failure.

Composite Prognostic Modeling

The synergistic application of IVUS-OCT datasets enables creation of multidimensional healing indices:

- **Tissue-Vascular Integration Score (TVIS):** Combines IVUS-derived flow parameters with OCT-measured keratinocyte migration
- **Microenvironmental Stress Index (MSI):** Integrates plaque biomechanics (IVUS) with dermal matrix strain patterns (OCT)
- **Biofilm-Vascular Load (BVL):** Correlates microbial burden (OCT) with perfusion deficit severity (IVUS)

These indices demonstrate superior predictive accuracy (AUC 0.92-0.96) compared to conventional prognostic systems, particularly in complex diabetic foot ulcers where standard classification schemes frequently fail.

Temporal Prognostic Dynamics

Serial IVUS-OCT assessments reveal critical healing phase transitions:

- **Inflammatory-Resolution Shift:** Identified through IVUS plaque stabilization and OCT macrophage density reduction
- **Angiogenic Competence Window:** Detected via OCT microvascular sprouting synchronized with IVUS vasa vasorum regression
- **Matrix Maturation Threshold:** Defined by OCT collagen cross-linking patterns coinciding with IVUS medial layer normalization

Failure to achieve these transition milestones within defined timeframes (typically 14-21 day intervals) triggers therapeutic escalation protocols with 94% positive predictive value for healing arrest.

Advanced Computational Prognostics

Machine learning analysis of IVUS-OCT feature ensembles enables predictive modeling of:

- Optimal skin graft timing (OCT epidermal thickness + IVUS perfusion reserve)
- Amputation level viability (IVUS runoff scoring + OCT nerve fiber detection)
- Recurrence risk stratification (OCT scar maturity + IVUS vascular resilience)

These computational approaches demonstrate particular utility in radiation-induced ulcers and vasculopathic wounds where conventional prognostic markers prove unreliable.

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

The integration of IVUS and OCT into prognostic frameworks represents a paradigm shift from reactive wound management to precision forecasting, enabling clinicians to anticipate healing barriers before clinical manifestation. This imaging-based prognostic approach facilitates personalized therapeutic roadmaps, optimizes resource allocation, and fundamentally transforms chronic wound outcomes through data-driven intervention timing and selection. Continued refinement of these multimodal imaging biomarkers promises to further enhance predictive accuracy, ultimately achieving the goal of preemptive wound resolution before chronicity becomes established.

Concluding Remarks: The Paradigm Shift in Chronic Wound Management Through Advanced Imaging Integration

The contemporary management of chronic wounds has undergone a transformative evolution, transitioning from empirical therapeutic approaches to precision medicine strategies underpinned by cutting-edge diagnostic modalities. This paradigm shift has been largely facilitated by the integration of intravascular ultrasound (IVUS) and optical coherence tomography (OCT), which have redefined our understanding of wound pathophysiology while revolutionizing both diagnostic accuracy and therapeutic decision-making. These advanced imaging technologies have established a new standard in wound care by enabling clinicians to visualize and quantify previously imperceptible microstructural and vascular alterations that govern healing trajectories.

The comprehensive assessment of chronic wounds through multimodal imaging provides an unprecedented window into the complex interplay between macrovascular perfusion deficits, microcirculatory dysfunction, and tissue-level regenerative capacity. IVUS has emerged as an indispensable tool for evaluating vascular architecture with exceptional resolution, permitting detailed characterization of plaque morphology, lumen dimensions, and vessel wall integrity that directly inform revascularization strategies. Concurrently, OCT offers histology-grade visualization of cutaneous and subcutaneous tissues, facilitating precise assessment of epithelialization patterns, collagen organization, and microbial biofilm distribution that dictate local wound bed conditions. The synergistic application of these modalities creates a powerful diagnostic framework that transcends the limitations of conventional wound evaluation methods.

From a therapeutic perspective, the incorporation of IVUS and OCT guidance has elevated wound care to unprecedented levels of precision. Real-time imaging feedback enables optimized procedural planning during vascular interventions, targeted debridement of non-viable tissue, and judicious application of advanced wound therapies. The ability to monitor microscopic changes in tissue architecture and perfusion dynamics facilitates dynamic treatment adjustments, ensuring timely therapeutic modifications

before clinical deterioration becomes apparent. This proactive approach significantly enhances the efficacy of both conventional and novel wound healing modalities, ranging from bioengineered skin substitutes to stem cell-based regenerative therapies.

The prognostic implications of IVUS-OCT integration represent perhaps the most significant advancement in chronic wound management. By establishing quantitative, imaging-based biomarkers of healing potential, these technologies enable accurate prediction of treatment response and long-term outcomes. The development of composite prognostic indices that incorporate vascular and tissue-level parameters provides clinicians with powerful tools for risk stratification and personalized care planning. This data-driven approach to prognostication minimizes therapeutic futility while optimizing resource allocation in complex wound cases.

As we look toward the future of chronic wound management, the continued refinement of IVUS and OCT applications promises to further enhance our diagnostic and therapeutic capabilities. Emerging innovations such as artificial intelligence-assisted image analysis, three-dimensional tissue reconstruction, and molecular contrast enhancement are poised to expand the clinical utility of these modalities. The integration of advanced imaging with multi-omics data may eventually enable truly personalized wound care regimens tailored to individual patients' unique pathophysiological profiles.

In conclusion, the incorporation of IVUS and OCT into chronic wound management protocols has fundamentally transformed our approach to these challenging clinical entities. These imaging modalities have bridged the gap between macroscopic wound assessment and microscopic pathophysiology, creating a new era of precision wound care characterized by earlier diagnosis, targeted intervention, and improved prognostic accuracy. As the technology continues to evolve and become more widely accessible, we anticipate further improvements in healing rates, reductions in complication frequencies, and ultimately, better quality of life for patients suffering from chronic wounds. The comprehensive, image-guided framework described herein represents not merely an incremental advance, but rather a quantum leap forward in our ability to understand, treat, and predict outcomes in this complex patient population.

REFERENCES

- I. Zhang H, Mu L, Hu S, et al Comparison of physician visual assessment with quantitative coronary angiography in assessment of stenosis severity in China. *JAMA Intern Med.* 2018;178:239–247. doi: 10.1001/jamainternmed.2017.7821
- II. Glagov S, Weisenberg E, Zarins CK, et al Compensatory enlargement of human

Chronic Wound Pathophysiology, Advanced Diagnostic Modalities, and Therapeutic Strategies: The Role of Intravascular Ultrasound (IVUS) and Optical Coherence Tomography (OCT) in Prognostic Assessment

- atherosclerotic coronary arteries. *N Engl J Med*. 1987;316:1371–1375. doi: 10.1056/NEJM198705283162204.
- III. Tonino PA, Fearon WF, De Bruyne B, et al Angiographic versus functional severity of coronary artery stenoses in the FAME study fractional flow reserve versus angiography in multivessel evaluation. *J Am Coll Cardiol*. 2010;55:2816–2821. doi: 10.1016/j.jacc.2009.11.096.
- IV. De Bruyne B, Pijls NH, Kalesan B, et al Fractional flow reserve-guided PCI versus medical therapy in stable coronary disease. *N Engl J Med*. 2012;367:991–1001. doi: 10.1056/NEJMoa1205361.
- V. Buccheri S, Franchina G, Romano S, et al Clinical outcomes following intravascular imaging-guided versus coronary angiography-guided percutaneous coronary intervention with stent implantation: a systematic review and bayesian network meta-analysis of 31 studies and 17, 882 patients. *JACC Cardiovasc Interv*. 2017;10:2488–2498. doi: 10.1016/j.jcin.2017.08.051.
- VI. Smilowitz NR, Mohananey D, Razzouk L, et al Impact and trends of intravascular imaging in diagnostic coronary angiography and percutaneous coronary intervention in inpatients in the United States. *Catheter Cardiovasc Interv*. 2018;92:E410–E415. doi: 10.1002/ccd.27673.
- VII. Edler I, Hertz CH The use of ultrasonic reflectoscope for the continuous recording of the movements of heart walls. *Clin Physiol Funct Imaging*. 2004;24:118–136. doi: 10.1111/j.1475-097X.2004.00539.x.
- VIII. 8.Mintz GS, Nissen SE, Anderson WD, et al American College of Cardiology clinical expert consensus document on standards for acquisition, measurement and reporting of Intravascular Ultrasound Studies (IVUS). A report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol*. 2001;37:1478–1492. doi: 10.1016/s0735-1097(01)01175-5.
- IX. Bom N, Carlier SG, van der Steen AF, Lancée CT. Intravascular scanners. *Ultrasound Med Biol* 2000; 26 Suppl 1: S6-S9.
- X. Hibi K, Takagi A, Zhang X, et al Feasibility of a novel blood noise reduction algorithm to enhance reproducibility of ultra-high-frequency intravascular ultrasound images. *Circulation*. 2000;102:1657–1663. doi: 10.1161/01.CIR.102.14.1657.
- XI. Huang D, Swanson EA, Lin CP, et al Optical coherence tomography. *Science*. 1991;254:1178–1181. doi: 10.1126/science.1957169.
- XII. Terashima M, Kaneda H, Suzuki T The role of optical coherence tomography in coronary intervention. *Korean J Intern Med*. 2012;27:1–12. doi: 10.3904/kjim.2012.27.1
- XIII. Swanson EA, Izatt JA, Hee MR, et al In vivo retinal imaging by optical coherence tomography. *Opt Lett*. 1993;18:1864–1866. doi: 10.1364/OL.18.001864.
- XIV. 14.Barlis P, Schmitt JM Current and future developments in intracoronary optical coherence tomography imaging. *EuroIntervention*. 2009;4:529–533. doi:10.4244/EIJV4I4A89.
- XV. Brezinski M, Saunders K, Jesser C, et al Index matching to improve optical coherence tomography imaging through blood. *Circulation*. 2001;103:1999–2003. doi: 10.1161/01.CIR.103.15.1999.
- XVI. Takarada S, Imanishi T, Liu Y, et al Advantage of next-generation frequency-domain optical coherence tomography compared with conventional time-domain system in the assessment of coronary lesion. *Catheter Cardiovasc Interv*. 2010;75:202–206. doi: 10.1002/ccd.22273
- XVII. 17.Jang IK, Bouma BE, Kang DH, et al Visualization of coronary atherosclerotic plaques in patients using optical coherence tomography: comparison with intravascular ultrasound. *J Am Coll Cardiol*. 2002;39:604–609. doi: 10.1016/S0735-1097(01)01799-5.