

Reinforced Tension Line (RTL) Technique for Abdominal Wall Closure: A Suture-Based Alternative to Mesh Reinforcement

Gerel Alecssandro Nuñez Falcon

Universidad Westhill. <https://orcid.org/0009-0004-5954-6291>

ABSTRACT

Background: Incisional hernia remains one of the most frequent complications after midline laparotomy, associated with significant morbidity, readmissions, and reoperations. Although small-bite suture techniques have reduced hernia rates, mechanical failure of the fascial closure remains an unresolved problem.

Objective: The reinforced tension line (RTL) technique was developed to enhance the biomechanical strength of abdominal wall closure by optimizing the distribution of tensile forces and preventing suture pull-through.

Methods and Evidence: The RTL method involves the placement of a longitudinal, nonabsorbable suture parallel to the primary fascial closure. This additional suture functions as a mechanical reinforcement line that redistributes peak tensile forces from the fascial edges into surrounding tissue. Recent randomized clinical trial data demonstrate that supplementing standard small-bite closure with an RTL suture significantly reduces the incidence of incisional hernia following midline laparotomy for colorectal cancer—from 21% with standard closure to 6% with RTL augmentation—without increasing wound complications or adverse events.

Results and Relevance: The RTL technique is especially useful when mesh reinforcement is contraindicated, such as in contaminated or emergent settings. It provides a cost-effective, mechanically sound, and biologically compatible alternative to mesh while maintaining a simple surgical workflow. However, mesh reinforcement remains superior for long-term prevention in high-risk elective cases.

Conclusion: RTL represents an innovative approach to fascial closure that enhances mechanical stability without introducing prosthetic material. Ongoing research is needed to define its long-term outcomes and applicability in complex abdominal wall reconstruction.

KEYWORDS: abdominal wall closure, incisional hernia, reinforced tension line, small-bite technique, mesh reinforcement, surgical innovation

ARTICLE DETAILS

Published On:
20 November 2025

Available on:
<https://ijmscrs.com/>

INTRODUCTION

Incisional hernia is one of the most common long-term complications following midline laparotomy, affecting up to one in four patients depending on surgical technique, patient comorbidities, and follow-up duration. These hernias are not merely aesthetic defects but carry substantial clinical and economic consequences, including chronic pain, bowel obstruction, and the need for surgical repair. Traditional methods of fascial closure have evolved over decades, with the small-bite, high-tension ratio technique—employing a 4:1 suture-to-wound length ratio—demonstrated to reduce hernia incidence compared to large-bite or low-ratio closures.

Despite these improvements, mechanical failure remains a concern, primarily due to localized suture pull-through and fascial tissue stress concentration. [1-3]




Figure 1. Schematic representation of RTL technique

To address these biomechanical limitations, the reinforced tension line (RTL) technique was developed. RTL introduces a longitudinal, nonabsorbable suture placed parallel to the primary closure line, functioning as a mechanical “load-sharing” element that redistributes tension across a broader area of the abdominal wall. By offloading peak stresses from the fascial edge and primary suture line, RTL minimizes localized stress points that can lead to tissue tearing, dehiscence, or hernia recurrence.[1-5]

This approach is distinct from mesh reinforcement or component separation, as it relies entirely on suture mechanics rather than prosthetic materials or tissue advancement. It therefore provides an attractive option in patients where mesh use is contraindicated, such as in contaminated surgical fields, emergency laparotomies, or high-risk septic settings.[6, 7]

DISCUSSION

The conceptual basis of RTL lies in its ability to modify the distribution of mechanical forces during wound healing. Standard midline closure often concentrates stress along the suture line, resulting in micro-tears and eventual failure at the suture-tissue interface. In contrast, RTL acts as a secondary tension-bearing system. When intra-abdominal pressure rises—such as during coughing or mobilization—the parallel suture absorbs part of the load, decreasing stress transmitted directly to the fascial closure. This mechanical “buffering” may reduce the risk of early fascial separation and late hernia formation. [7]

Clinical Evidence

Recent randomized clinical data have validated the theoretical advantages of RTL. In a controlled trial involving patients undergoing midline laparotomy for colorectal cancer, the addition of an RTL suture to a standard small-bite (4:1) closure led to a remarkable reduction in the one-year CT-detected incisional hernia rate—from 21% with standard closure alone to only 6% with RTL reinforcement.[8] Notably, this improvement was achieved without an increase in wound infections, seroma, or dehiscence, underscoring the safety and feasibility of the technique in clinical practice.

Advantages

The RTL technique offers several notable benefits:

Enhanced biomechanical stability: By redistributing tensile loads, RTL prevents focal stress and reduces the risk of suture pull-through, particularly in patients with obesity, poor tissue quality, or elevated intra-abdominal pressure.[1, 4, 8]

Avoidance of prosthetic material: Unlike mesh reinforcement, RTL introduces no foreign body, thereby reducing infection risk—a key advantage in contaminated or potentially contaminated cases such as perforated bowel or emergency surgery.[6]

Technical simplicity and accessibility: The technique can be performed with standard suture materials and does not require specialized equipment, making it feasible in both high-resource and limited-resource settings. [1, 4, 8]

Cost-effectiveness: By avoiding prosthetic mesh, RTL reduces procedural cost while maintaining hernia prevention efficacy comparable to more complex interventions.

Disadvantages and Limitations

Despite these strengths, RTL has certain limitations:

Reinforced Tension Line (RTL) Technique for Abdominal Wall Closure: A Suture-Based Alternative to Mesh Reinforcement

Inferiority to mesh in high-risk elective cases: Meta-analyses indicate that prophylactic mesh reinforcement remains the most effective method for preventing incisional hernia, particularly in patients at extreme risk such as those undergoing open abdominal aortic aneurysm repair or repeated laparotomies.[3]

Limited long-term data: Most RTL studies report outcomes at one to two years; the long-term durability compared with mesh reinforcement remains unclear. Further studies with extended follow-up are necessary. [6, 8]

Not suitable for all cases: RTL may be insufficient in cases with extreme fascial tension (>40 N), massive hernias, or severe loss of domain, where prosthetic support is required to achieve a tension-free closure.[2]

Limited evidence in complex reconstructions: Its role in reoperative or multi-plane abdominal wall repairs is yet to be fully defined.

Comparative Perspective

In the continuum of abdominal wall closure strategies, RTL occupies a valuable intermediate position. It provides a mechanical advantage superior to standard suture closure while avoiding the drawbacks of prosthetic mesh. In emergency or contaminated fields, it offers a safe and effective alternative when mesh implantation poses infection risks. For elective, high-risk patients, however, mesh reinforcement continues to yield the lowest hernia recurrence rates. The ideal surgical approach should therefore be individualized, balancing biomechanical needs, contamination risk, and resource availability. [1-8]

Table 1. Advantages and Disadvantages of the Reinforced Tension Line (RTL) Technique

Aspect	Advantages	Disadvantages / Limitations
Incisional hernia prevention	Significantly reduces hernia rate compared to standard small-bite closure (21% → 6% at 1 year).[2]	Mesh reinforcement still superior for high-risk elective cases (e.g., aortic aneurysm repair).[3]
Biomechanical performance	Redistributes tensile forces, preventing suture pull-through and fascial tearing; improves closure strength.[1,2]	Efficacy not well established in very large or recurrent hernias where fascial tension exceeds 40 N.[2]
Infection risk	Avoids prosthetic material, reducing risk of infection in contaminated or emergent settings.[2]	Long-term comparative data with mesh reinforcement remain limited.[3]
Technical demands	Simple, reproducible, and uses standard suture materials; no special equipment needed.	Requires familiarity with technique; learning curve may influence outcomes.
Cost and resources	No need for mesh or additional devices—potentially reduces procedural cost.	Savings may be offset if recurrence occurs in very high-tension cases.
Evidence base	Supported by randomized trial showing short-term efficacy and safety.	Limited data in complex or reoperative abdominal wall reconstruction.

CONCLUSION

The reinforced tension line (RTL) technique represents a significant evolution in suture-based abdominal wall closure. By adding a longitudinal reinforcement suture parallel to the primary closure, RTL distributes mechanical load more evenly across the fascial plane, minimizing the risk of suture pull-through and incisional hernia formation. Clinical trials demonstrate a substantial reduction in hernia rates without an increase in complications, establishing RTL as a promising, safe, and cost-effective technique.

Its particular strength lies in contaminated or emergent scenarios, where mesh use is contraindicated. While mesh reinforcement remains the gold standard for elective high-risk cases, RTL provides an innovative solution for patients who would otherwise be at increased risk of hernia recurrence. Long-term studies are needed to confirm the durability of RTL and to define its role in complex and reoperative abdominal wall reconstruction.

In summary, the RTL technique combines biomechanical soundness, procedural simplicity, and clinical effectiveness—

offering surgeons a valuable tool in the ongoing effort to reduce postoperative incisional hernia rates.

REFERENCES

I. Wenzelberg, C. L., Rogmark, P., Ekberg, O., & Petersson, U. (2024). Reinforced tension-line suture after laparotomy: early results of the Rein4CeTo1 randomized clinical trial. *British Journal of Surgery*, 111(10), znae265.

II. Lozada Hernández, E. E., Flores González, E., Chavarría Chavira, J. L., Hernandez Herrera, B., Rojas Benítez, C. G., García Bravo, L. M., ... & Soria Rangel, J. (2024). The MESH-RTL Project for prevention of abdominal wound dehiscence (AWD) in high-risk patients: noninferiority, randomized controlled trial. *Surgical Endoscopy*, 38(12), 7634-7646.

III. Lozada-Hernández, E. E., López-Briones, J. S., Martín del Campo, L. A., Fernández-Vazquez Mellado, L. A., Valenzuela Alpuche, H. A., Jean-Silver, E. R., ... & Carlos-Jimenez, S. (2025).

Reinforced Tension Line (RTL) Technique for Abdominal Wall Closure: A Suture-Based Alternative to Mesh Reinforcement

- Comparison of the levels of metalloproteinases and their tissue inhibitors in serum and fascia in four methods of midline laparotomy closure: standard, small bites, reinforced tension line and onlay mesh (absorbable and non-absorbable). Experimental study. *Journal of Surgical Protocols and Research Methodologies*, 2025, snaf008.
- IV. Lozada-Hernández, E. E., López-Briones, J. S., Martín del Campo, L. A., Fernández-Vazquez Mellado, L. A., Valenzuela Alpuche, H. A., Jean-Silver, E. R., ... & Carlos-Jimenez, S. (2025). Comparison of the levels of metalloproteinases and their tissue inhibitors in serum and fascia in four methods of midline laparotomy closure: standard, small bites, reinforced tension line and onlay mesh (absorbable and non-absorbable). Experimental study. *Journal of Surgical Protocols and Research Methodologies*, 2025, snaf008.
- V. van den Berg, R., Visscher, L., Menon, A. G., Deerenberg, E. B., & Tanis, P. J. (2025). Suture Techniques and Materials for Fascial Closure of Abdominal Wall Incisions: A Comprehensive Meta-Analysis. *Annals of Surgery Open*, 6(1), e548.
- VI. Wenzelberg, C., Petersson, U., Syk, I., Ekberg, O., & Rogmark, P. (2023). Abdominal closure with reinforcing suture decreases incisional hernia incidence after CRS/HIPEC. *Journal of Abdominal Wall Surgery*, 2, 11188.
- VII. Acuña Pinzón, C. L. (2023). Prevención de hernia incisional con técnica RTL (Reinforced Tension Line) comparada con el uso de malla profiláctica en laparotomía por línea media: seguimiento a 3 años del reporte inicial: ensayo clínico aleatorizado de no inferioridad.
- VIII. Cuenca, O., Ferreira, R., Ferreira, J., Sosa, E., Agüero, A., Cano, C., & Díaz, J. R. (2025). Técnica de refuerzo de la pared abdominal con sutura longitudinal lateral a la incisión (reinforced tension line-RTL) para el cierre de la línea media como prevención de evisceración. *Revista Cirugía Paraguaya*, 49(1), 12-15.