

Perforator Flaps: Evolution, Classification, and Clinical Selection Algorithms

Diego Clemente Moreno¹, Sergio Jacobo Flemate², Clyvia Coral Medina Lavenant³, Víctor Esteban Kuri⁴, Alexis Esquivel Estrada⁴, Joselyn Ugalde Rivera⁵, Dylan Ruiz Reyes⁶, Rogelio Romero Vázquez⁷

¹IMSSS Hospital General Regional no. 2 El Marqués, Orcid: 0009-003-1784-7486

²Hospital General Reynosa.Orcid : 0009-0000-9161-753X

³Hospital General de Zona 16 IMSS, Orcid: 0009-0004-8230-141X

⁴Centro Medico Licenciado Adolfo López Mateos Orcid: 0009-0002-3056-8353

⁵Hospital General de Querétaro, 0009-0006-5710-6190

⁶Hospital General de México Dr. Eduardo Liceaga, Orcid: 0009-0000-1553-0379

⁷IMSS HGR No.270, Orcid: 0009-0001-6887-982X

INTRODUCTION

Modern reconstructive surgery is defined by the constant pursuit of maximum efficacy with minimal donor site morbidity. In this context, **Perforator Flaps** have become the gold standard in numerous reconstructive situations, representing the highest echelon of surgical refinement [1]. These flaps allow the transfer of fasciocutaneous tissue, and sometimes composite tissue, based exclusively on a blood vessel that passes through a deep plane (muscle or septum)

before supplying the skin, thereby preserving the functional integrity of the underlying muscle [2]. The objective of this review article is to provide plastic surgery residents with an updated and comprehensive reference that synthesizes the historical evolution of this technique, its systematic classification, and the essential clinical selection algorithms for its successful application in contemporary reconstructive practice.

Thematic Development

1. Historical Evolution: From Random Flap to Perforator Flap

The history of perforator flaps is a chronicle of the progressive understanding of cutaneous vascular anatomy.

Era	Dominant Concept	Key Milestones	Impact on Practice
Mid-20th Century	Random Cutaneous Flaps	Based on the subdermal plexus. Viability limited by the length-to-base ratio (maximum 2:1) [1].	Local and limited reconstructions.
1960s and 1970s	Axial and Musculocutaneous Flaps	Daniel and Taylor describe flaps based on the axial axis. Popularization of musculocutaneous flaps (Mathes and Nahai) [1].	Greater vascular reliability, but high donor site morbidity (muscle sacrifice) and suboptimal aesthetic results (bulky flaps).
1981	Fasciocutaneous Flaps	Ponten postulates the inclusion of fascia in the flap to improve viability ("superflaps") [1].	Reduction of muscular morbidity, but still lacking the current refinement.
1989	Birth of the Perforator Flap	Koshima and Soeda introduce the concept of the <i>Perforator Flap</i> based on a perforator from the inferior epigastric artery [1].	Fundamental milestone: Demonstration that tissue can be transferred with high vascular reliability without sacrificing the muscle.

Perforator Flaps: Evolution, Classification, and Clinical Selection Algorithms

Era	Dominant Concept	Key Milestones	Impact on Practice
1994 - Present	Modern Era	Blondeel and Boeckx popularize the DIEP (Deep Inferior Epigastric Perforator) flap for breast reconstruction [1].	Gold standard in reconstructive microsurgery, minimizing donor site morbidity.

The concept of the **angiosome** , described by Taylor in 1988, provided the physiological basis for the development of these flaps, by demonstrating that the skin and underlying tissues are supplied by interconnected three-dimensional vascular units [2].

2. CLASSIFICATION OF PERFORATOR FLAPS

The classification of perforator flaps is based on the anatomy of the vessel that passes through the deep plane to reach the skin. Understanding this classification is fundamental for dissection and nomenclature.

2.1. Classification of Perforator Vessels

Perforator vessels are classified according to the plane they traverse before reaching the fasciocutaneous tissue [2]:

Perforator Type	Anatomical Course	Clinical Implication
Musculocutaneous (MCp)	Passes through the muscle before perforating the fascia and reaching the skin.	Requires intramuscular dissection (separating muscle fibers) to isolate the pedicle. E.g.: DIEP Flap.
Septocutaneous (SCp)	Passes through the intermuscular septum before perforating the fascia and reaching the skin.	Dissection is simpler, as it does not require separation of muscle fibers. E.g.: ALT Flap (in some cases).
Direct Cutaneous (DCp)	Passes only through the fascia (not the muscle or septum).	Dissection is the simplest, as the pedicle is suprafascial.

2.2. Nomenclature and Clinical Examples

The modern nomenclature of perforator flaps combines the name of the source vessel with the abbreviation "P" for Perforator [2].

Flap	Source Vessel	Primary Clinical Application
DIEP	Deep Inferior Epigastric Artery	Breast reconstruction, trunk defects.
ALT	Anterolateral Thigh (Lateral Circumflex Femoral Artery)	Lower extremity, head, and neck defects.
TDAP	Thoracodorsal Artery	Partial breast reconstruction, trunk defects.
SGAP/IGAP	Superior/Inferior Gluteal Arteries	Gluteal reconstruction, sacral defects.
Propeller Flaps	Local Perforators (e.g., Posterior Tibial Artery)	Small/medium defects in extremities, especially distal ones.

3. Clinical Selection Algorithms and Preoperative Planning



Figure 1.-Lumbar defect.




Figure 2.-DIEP flap Marking.

The selection of the optimal perforator flap is an algorithmic process that balances the needs of the defect with the availability and morbidity of the donor site.

3.1. Decision Criteria

The decision-making process is based on a comprehensive evaluation of three pillars [3]:

1 Defect Characteristics:

◦ **Size and Location:** Large or complex defects often require free flaps (DIEP, ALT). Small/medium defects in extremities can be solved with local propeller flaps.

◦ **Tissue Composition:** Determine whether only skin and fat are required (DIEP flap) or if composite tissue (fascia, tendon

2 Patient Characteristics:

◦ **Comorbidities:** The patient's vascular status is crucial.
◦ **Donor Site Availability:** Evaluate tissue quality, presence of previous scars, and suitability of recipient vessels.

3 Reconstructive Goal:

◦ **Functionality and Aesthetics:** The main benefit of perforator flaps is **minimal donor site morbidity** (muscle preservation), which translates into better functional and aesthetic recovery.

3.2. Preoperative Planning and Perforator Mapping

The key to success in perforator flap surgery lies in precise preoperative planning. The goal is to identify the **ideal perforator** that meets three criteria [3]:

- 4 **Adequate Caliber and Pulsatility:** Sufficient to perfuse the entire flap.
- 5 **Central Location:** To ensure homogeneous perfusion of the tissue to be transferred.
- 6 **Minimal Intramuscular Course:** To simplify dissection and reduce muscle damage.



Figure 2.-SGAP bilateral flap.




Figure 4.-DIEP Flap.

Various tools are used for perforator mapping:

Mapping Method	Advantages	Disadvantages	Clinical Role
Ultrasound Doppler	Easy to use, portable, low cost.	Low specificity, subjective, high risk of false positives.	Initial orientation and quick marking.
Color Duplex Doppler	High sensitivity, accurately evaluates caliber and flow.	Long examination time, highly technician-dependent.	Confirmation of doubtful perforators.
Multidetector Computed Tomography (MDCT) / CT Angiography	High sensitivity and specificity (100% predictive value), 3D reconstruction, precise coordinate mapping [3].	Requires advanced technology and exposure to radiation/contrast.	Method of choice for complex free flaps (e.g., DIEP).

Super Thin Flaps

The evolution of perforator flaps does not stop at muscle preservation. The most recent trend in reconstructive microsurgery focuses on the refinement of the thickness of the transferred tissue, giving rise to the concept of Super Thin Flaps or Thinned Perforator Flaps. This technique represents the pinnacle of refinement, seeking aesthetic and functional results that completely mimic the surrounding tissue [4].

Principles and Justification

Thinning Techniques.

The thinning of a perforator flap can be performed at two key moments, which influences the complexity and the result:

Technique	Description	Advantages	Disadvantages
Primary Thinned Elevation	The flap is intentionally raised in a superficial plane (suprafascial or even above the superficial subcutaneous fascia) from the beginning of the dissection [4].	Single-stage procedure. Maximum tissue malleability.	Higher risk of vascular compromise if the perforator is damaged or if perfusion is marginal.
Secondary Thinning	The flap is thinned in a subsequent surgery, once the flap has taken and its vascularization is established.	Greater vascular safety. Allows for more precise contouring.	Requires a second surgical procedure.
Combined Techniques	Use of assisted liposuction or preoperative delay techniques to improve flap viability before thinning [4].	Increases safety in the primary thinning of large flaps (e.g., DIEP) [4].	Greater complexity and total surgical time.

Advanced Preoperative Planning

The safety of a super thin flap directly depends on the precise identification of the perforator and its superficial vascular network. Imaging tools have evolved to support this need:

Mapping Method	Specific Contribution for Super Thin Flaps
CT Angiography (MDCT)	Remains the gold standard for mapping the perforator's course from the source vessel to the fascia. Crucial for planning the dissection [5].
Ultra-High-Frequency Ultrasound (UHF-US)	Allows detailed visualization of the superficial vascular anatomy and tissue thickness, facilitating the planning of the elevation in the suprafascial or superficial subcutaneous plane [6].

Recent advance in the field of super thin flaps is the reappraisal of the superficial circumflex iliac artery perforator (SCIP) flap. Traditionally limited by a short pedicle, a new technique proposes basing the flap on the perforators of the inferolateral branches of the SCIA. This approach offers superior anatomical consistency and allows, through simplified retrograde dissection, pedicle lengthening to include the entire SCIA with minimal muscle dissection. The most notable feature is that these perforators extend directly to the dermal plexus, facilitating the primary elevation of a super thin flap without the need for microsurgical defatting. This flap is ideal for the reconstruction of superficial soft-tissue defects that require fine contouring and minimal volume transfer. [8].

Clinical Applications

The super thin flap has become the preferred option for:

The main objective of super thin flaps is to avoid the undesirable "patch effect" that occurs when a flap is too bulky for the defect to be covered. This is especially critical in areas of delicate contour such as the face, neck, hands, and feet. The historical progression is summarized as:

- 1.Musculocutaneous Flaps: High donor site morbidity.
- 2.Perforator Flaps: Muscle preservation, reduced morbidity.
- 3.Super Thin Flaps: Thickness refinement, aesthetic optimization [4].

- Extremity Reconstruction: The thinned anterolateral thigh (ALT) flap is widely used for defects in the leg and foot, where excessive bulk is a functional and aesthetic problem [6].
- Facial and Neck Reconstruction: Thinness is essential for aesthetic integration in these areas of fine contour.
- Breast Reconstruction: The thinning of the DIEP flap, whether primary or secondary, is a common approach to optimize the contour and shape of the reconstructed breast [7].

CONCLUSIONS

Perforator flaps represent the culmination of evolution in reconstructive surgery, offering a solution that combines the vascular reliability of musculocutaneous flaps with minimal donor site morbidity. One must master not only the surgical technique but also a deep understanding of vascular anatomy

Perforator Flaps: Evolution, Classification, and Clinical Selection Algorithms

(MCp, SCp, DCp classification) and the rigorous application of clinical selection algorithms.

Preoperative planning, especially with the use of CT Angiography (MDCT), is indispensable for success, allowing for the selection of the ideal perforator and the preservation of muscle function. By applying these principles, the surgeon in training will be equipped to make informed reconstructive decisions, significantly improving functional and aesthetic outcomes for their patients.

REFERENCES

- I. Redalyc. Colgajos de perforantes. Principios Básicos y Aplicaciones Clínicas. *Revista Med*, 2012.:
<https://www.redalyc.org/pdf/910/91025872005.pdf>
- II. Binder, J.P., Servant, J.M., Revol, M. Colgajos perforantes. *EMC - Cirugía Plástica Reparadora y Estética*, 2012.
<https://www.sciencedirect.com/science/article/abs/pii/S1634214312627472>
- III. Masià, J. Planificación preoperatoria de los colgajos de perforantes. *Cirugía Plástica Ibero-Latinoamericana*, 2006.
http://scielo.isciii.es/scielo.php?script=sci_arttext&pid=S0376-78922006000400003
- IV. Franchi, A. Deep Inferior Epigastric Perforator Flap Preconditioning... *Plastic and Reconstructive Surgery*, 2025.
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11932617/>
- V. [5] Chim, H. P135. SUPERTHIN PROFUNDA ARTERY PERFORATOR... *Plastic and Reconstructive Surgery*, 2022.
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8984374/>
- VI. [6] Visconti, G. Thin and superthin perforator flap elevation based on... *Plastic and Reconstructive Surgery*, 2020.
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7398805/>
- VII. [7] Zhou, J. Super-thin anterolateral thigh flap for reconstruction of the... *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 2023.
<https://journals.sagepub.com/doi/10.1177/10225536231181706>
- VIII. [8] Zhang Y, Zeng A. An Easy Way to Harvest a Superthin SCIP Flap with Long Pedicle: Reappraisal of the Inferolateral Branches of the SCIA. *Plast Reconstr Surg*. 2023 Nov 1;152(5):1100-1104. doi: 10.1097/PRS.00000000000010338. Epub 2023 Mar 3. PMID: 36862953.