

Redefining Breast Rejuvenation: Current Aesthetic Approaches in Mastopexy

Axel Roberto Rosas Aleman¹, Jazmyn Elizabeth Rodríguez Contreras², Estefany Marlen Barragán Jiménez², Jesús Insúa Pohls³, Félix Steven Bojórquez Mora¹, José Noé Corrales Medina¹, Iván Humberto Acevedo Ojeda⁴, Francisco Javier Cano Palacios⁵

¹Universidad Autónoma de Sinaloa, Culiacán, Sinaloa, Mexico

²Hospital General de Zona No. 3, IMSS, Aguascalientes, Mexico

³Hospital General Regional 1, Ciudad Obregón, Sonora, Mexico

⁴Universidad Marista de Mérida, Mérida, Yucatán, Mexico

⁵Hospital Regional Dr. Manuel Cárdenas de la Vega, ISSSTE, Mexico

ABSTRACT

Background: Mastopexy plays a central role in aesthetic breast rejuvenation, targeting correction of ptosis while restoring youthful projection, upper pole fullness, and symmetry. Recent innovations emphasize optimizing breast shape and minimizing scar burden without compromising long-term outcomes.

Methods: This review summarizes current concepts and evidence-based strategies in mastopexy, focusing on patient selection, technique planning, parenchymal reshaping, and approaches to reduce recurrent ptosis. Special attention is given to vertical and limited-scar patterns, augmentation–mastopexy considerations, and emerging internal support technologies.

Results: Contemporary mastopexy techniques demonstrate high rates of patient satisfaction and low major complication rates when performed by experienced surgeons. Aesthetic shortcomings tend to relate to scar quality and recurrent ptosis rather than functional failures. Hybrid approaches integrating implants, fat grafting, and soft-tissue support systems show promise in improving upper pole stability and aesthetic longevity.

Conclusion: Aesthetic mastopexy continues to evolve toward personalized, anatomy-driven strategies that enhance shape retention and optimize scar placement. Future development of internal support and hybrid volume restoration may further refine aesthetic durability.

KEYWORDS: Mastopexy, Breast ptosis, Aesthetic breast surgery

ARTICLE DETAILS

Published On:
30 December 2025

Available on:
<https://ijmscrs.com/>

Mastopexy is a cornerstone of aesthetic breast surgery aimed at correcting ptosis while optimizing breast shape, upper pole fullness, and scar quality. Modern approaches emphasize harmonizing breast volume and position with the thoracic frame, prioritizing long-term projection and stable upper pole contour over simple elevation of the nipple–areola complex (NAC) [1].

Aesthetic Goals and Patient Selection

Ideal candidates include those with breast ptosis and loss of upper pole fullness who are satisfied with their breast volume or seek modest augmentations through augmentation–mastopexy. Matching ptosis severity to the least-invasive technique that achieves acceptable nipple position and shape is critical for success and patient satisfaction [1]




Figure 1. Preoperative planning and postoperative appearance




Figure 2. Follow up with nice results

Preoperative planning evaluates:

- NAC position relative to the IMF
- Skin quality and parenchymal distribution
- Chest wall anatomy and asymmetry
- Patient tolerance to scar burden

Technique Selection and Scar Strategy

Limited-scar and vertical approaches have become widely adopted for their ability to limit scar burden while maintaining or improving breast shape. Vertical mastopexy, described by Hall-Findlay, emphasizes conical reshaping and upper pole projection with a vertical scar and short horizontal limb when necessary [2].

Technique considerations:

- Periareolar mastopexy — for mild ptosis; minimal scars but risk of areolar widening
- Vertical mastopexy — good for moderate ptosis; high projection, concealed scars
- Wise-pattern mastopexy — powerful reshaping for severe ptosis but longer scars

A systematic review confirms mastopexy as a reliable and satisfying aesthetic procedure, with most postoperative dissatisfaction relating to scarring or recurrence of ptosis rather than major complications [3].

Outcomes and Patient Satisfaction

Complication rates remain low in experienced hands. Minor issues include scarring irregularities, delayed healing, and lower pole elongation. Revisions often address cosmetic refinements rather than functional failures.

Stevens et al. and others report high patient satisfaction (>85%) with overall shape and nipple position after mastopexy procedures [4].

Augmentation–Mastopexy and Upper Pole Aesthetics

For patients with volume deficiency and ptosis, combining implant augmentation with mastopexy can restore upper pole fullness and projection. However, skin envelope tightening with implant insertion increases mechanical demands on tissues. Spear et al. demonstrated high satisfaction rates but noted higher reoperation rates compared to staged procedures due to recurrent ptosis or implant issues [5].

A large meta-analysis by Khavanin et al. confirmed that single-stage augmentation–mastopexy is safe when patient selection is appropriate [6].

Contemporary Trends: Mesh, Fat Grafting, and Hybrid Techniques

Current aesthetic goals extend beyond elevating the NAC to improving three-dimensional contour and long-term soft-tissue support. Innovations include:

- Internal support systems (biologic/synthetic mesh)
- Hybrid augmentation combining implants with fat grafting to improve contour and softness
- Parenchymal reshaping techniques that reduce dependence on skin for support

These strategies aim to reduce recurrent ptosis and maintain upper pole fullness over time.

Table 1. Comparison of Common Mastopexy Techniques with Aesthetic Focus

Technique	Ideal Indications	Key Advantages	Aesthetic Limitations	Typical Scars
<i>Periareolar (Circumareolar/Benelli)</i>	Mild ptosis, minor areolar repositioning, small breasts with good skin elasticity	Minimal scar, good for small adjustments, NAC symmetry correction	Risk of areolar widening, breast flattening, limited reshaping for moderate/severe ptosis	Circular scar around NAC
<i>Vertical Mastopexy (Hall-Findlay and variants)</i>	Moderate ptosis, desire for conical shape with projection	Enhances upper pole projection, long-term shape stability, limited scars	May require short horizontal scar in some cases, “bottoming out” in poor tissue quality	Periareolar + vertical
<i>Wise-Pattern (Inverted-T)</i>	Severe ptosis, significant skin excess, massive weight loss patients	Greatest power of reshaping and skin tightening, precise control of breast footprint and lower pole	Longer scar burden, higher risk of visible scarring; relies more on skin for support if no parenchymal shaping is used	Periareolar + vertical + IMF transverse
<i>Auto-augmentation Mastopexy (Parenchymal rearrangement)</i>	Volume loss without implants, postpartum changes	Improved upper pole fullness without implants, avoids implant-related risks	Limited volume improvement compared with augmentation, technique dependent	Depends on mastopexy pattern
<i>Augmentation–Mastopexy (Single-Stage)</i>	Ptosis + volume deficiency	Restores projection and upper pole fullness, comprehensive rejuvenation in one procedure	Higher reoperation risk: recurrent ptosis, implant malposition, capsular contracture	Depends on mastopexy type + implant

Redefining Breast Rejuvenation: Current Aesthetic Approaches in Mastopexy

Hybrid Mastopexy (Implant + Fat Grafting)	Desire for soft contour and implant camouflage	Natural feel, refined shape, can reduce implant size and tensions	Requires multiple fat grafting sessions in some cases	Depends on mastopexy type
Mesh-Supported Mastopexy ("Internal Bra")	Poor skin quality, recurrent ptosis risk	Additional lower-pole support, improved long-term projection	Cost, variable evidence, foreign-body concerns	Depends on mastopexy type

CONCLUSION

Mastopexy continues to evolve toward techniques prioritizing durable projection, high aesthetic quality, and minimal scar burden. Vertical and limited-scar approaches provide consistent results for most patients, while Wise-pattern techniques remain indispensable in severe ptosis and massive weight loss. Augmentation–mastopexy offers comprehensive rejuvenation for selected patients but requires careful planning to reduce revision risks. Advances in internal support and hybrid volume enhancement demonstrate promising improvements in long-term outcomes. A tailored, anatomy-based approach remains essential for achieving optimal aesthetic satisfaction.

REFERENCES

- I. Rohrich RJ, Gosman AA, Brown SA, Reece EM. The limited scar mastopexy: current concepts and approaches to correct breast ptosis. *Plast Reconstr Surg.* 2004;114(7):1759-1770.
- II. Hall-Findlay EJ. A simplified vertical reduction mammoplasty: shortening the learning curve. *Plast Reconstr Surg.* 1999;104(3):748-759.
- III. di Summa PG, Oranges CM, Watfa W, et al. Systematic review of outcomes and complications in nonimplant-based mastopexy surgery. *J Plast Reconstr Aesthet Surg.* 2019;72(2):243-272.
- IV. Stevens WG, Stoker DA, Freeman ME, et al. Mastopexy revisited: a review of 150 consecutive cases for complications and aesthetic results. *Aesthetic Surg J.* 2004;24(3):189-196.
- V. Spear SL, Pelletiere CV, Menon N. One-stage augmentation combined with mastopexy: aesthetic results and patient satisfaction. *Aesthetic Plast Surg.* 2004;28(5):259-267.
- VI. Khavanin N, Jordan SW, Rambachan A, et al. A systematic review of single-stage augmentation-mastopexy. *Plast Reconstr Surg.* 2014;134(5):922-931.