

Topographic and Neurovascular Determinants of Dental Implant Complications: An Anatomical Review for Surgical Planning

K. Mardaleishvili^{1*}, T. Baramidze²

¹Oncology Research Medical Center, Tbilisi, Georgia

²Department of Head and Neck Oncology, BAU International University, Batumi, Georgia

ABSTRACT

Dental implantology is performed within an anatomically compact region in which osseous, vascular, lymphatic, sinus, and neural structures lie in close proximity. The objective of this narrative review is to reorganize the anatomical content of the source manuscript into a journal-style article focused on topographic and neurovascular risk assessment. The review examines maxillary and mandibular anatomical zones that determine surgical safety, including the maxillary sinus, Schneiderian membrane, nasopalatine canal, thin buccal cortical plate, infraorbital region, inferior alveolar canal, mental foramen, lingual cortical plate, and floor-of-mouth vascular spaces. Particular attention is given to how anatomical variation contributes to sinus perforation, implant malposition, intraosseous bleeding, hematoma, thermal osteonecrosis, and trigeminal-branch injury. The article emphasizes CBCT-based planning, preservation of safety margins, atraumatic osteotomy, copious irrigation, and systematic postoperative neurosensory monitoring. A precise anatomical framework remains essential for reducing preventable complications and improving functional outcomes after dental implant placement.

KEYWORDS: Dental implants, Maxillary sinus, Inferior alveolar nerve, Computed tomography, Surgical planning, Complications

ARTICLE DETAILS

Published On:
26 August 2026

Available on:
<https://ijmscrs.com>

Maxilla Maxillary sinus / Schneiderian membrane, Nasopalatine and incisive canals, Infraorbital region, Thin buccal cortical plate	Mandible Inferior alveolar canal, Mental foramen and anterior loop, Lingual cortical plate, Sublingual and submental vessels
Operative Risk Sinus perforation, cortical fenestration, vascular injury, thermal osteonecrosis	Functional Risk Paresthesia, dysesthesia, neuropathic pain, hematoma, airway-threatening swelling

Figure 1. Topographic risk zones relevant to implant planning.

I. INTRODUCTION

The clinical success of dental implant placement depends not only on implant design and operative technique but also on the surgeon's ability to interpret the maxillofacial anatomical complex as a functional system. The head and neck region contains the eye, nose, oral cavity, maxilla, mandible, hard and soft palate, salivary glands, cervical vessels, cranial nerves, and lymphatic pathways within a limited anatomical

field. Therefore, small deviations in implant depth, angulation, or osteotomy trajectory may affect neurovascular safety, sinus integrity, bone vitality, and postoperative function.

This article presents an anatomical review designed for journals focused on oral surgery, implantology, and clinical anatomy. Its central thesis is that complications after implant placement are often predictable when the maxilla and

Topographic and Neurovascular Determinants of Dental Implant Complications: An Anatomical Review for Surgical Planning

mandible are analyzed through topographic risk zones rather than only through available bone height and width.




Figure 2. Blood Vessels of the Head and Neck - Atlas of Anatomy. Head and Neuroanatomy. Michael Schuenke. (2015)

II. METHODOLOGICAL APPROACH

This manuscript is structured as a narrative anatomical review derived from the author-provided Georgian source text and supported by current literature on CBCT-based implant planning, odontogenic infection complications, inferior alveolar nerve injury, and peri-implant disease. The focus is not on presenting new clinical cases or statistical data, but on synthesizing anatomical mechanisms into a practical surgical-planning framework.

III. MAXILLARY ANATOMY AND IMPLANT-RELATED RISK

A. Sinus Floor and Schneiderian Membrane

The posterior maxilla is one of the most clinically sensitive regions for implant placement because the maxillary sinus floor may lie close to the alveolar ridge, especially after tooth loss and alveolar resorption. Sinus membrane perforation may occur during osteotomy preparation, implant insertion, or sinus-lift procedures. The consequences include postoperative sinusitis, oroantral communication, inflammatory thickening of the sinus mucosa, or implant migration into the sinus cavity.

Prevention requires three-dimensional evaluation of residual bone height, sinus septa, membrane thickness, and the spatial relationship between the planned implant and the sinus floor. Implant length, diameter, and angulation should be selected in relation to the sinus boundary and prosthetic axis, not merely by available ridge height.

B. Nasopalatine and Incisive Canal Region

In the anterior maxilla, the nasopalatine canal and incisive canal may be injured when implants are placed near the central incisors. Trauma to this region may produce pain, altered sensation, bleeding, local inflammation, or implant instability if the canal is used as an unintended osteotomy pathway. Radiographic identification of canal size, morphology, and distance from the proposed implant site is therefore essential.

C. Buccal Cortical Plate and Esthetic-Zone Vulnerability

The maxillary buccal cortical plate is frequently thin, particularly in the anterior maxilla. A deficient buccal wall increases the risk of fenestration, dehiscence, soft-tissue recession, implant exposure, and esthetic failure. The selection of smaller-diameter implants, palatal positioning when prosthetically acceptable, staged augmentation, and careful flap management may reduce these risks.

D. Infraorbital and Pterygopalatine Region

The infraorbital nerve, posterior superior alveolar vessels, and pterygopalatine vascular branches form a complex regional network around the posterior and superior maxilla. Although infraorbital nerve injury during routine dental implant placement is uncommon, procedures near premolar regions, sinus-lift windows, or the pterygopalatine fossa require caution because local vascular and neural structures may contribute to bleeding, sensory disturbance, or infection spread.

IV. MANDIBULAR ANATOMY AND IMPLANT-RELATED RISK

A. Inferior Alveolar Canal

The inferior alveolar canal is the principal anatomical limitation in posterior mandibular implant placement. Injury may occur through direct drill penetration, implant compression, heat generation, postoperative edema, hematoma, or secondary infection. Clinical outcomes may include paresthesia, anesthesia, dysesthesia, or chronic neuropathic pain affecting the lower lip, chin, gingiva, or mandibular teeth. A minimum safety distance should be preserved between the implant apex and the canal, with CBCT used to evaluate canal position, cortical boundaries, and anatomical variation.




Figure 3. Mandibular canal and inferior alveolar neurovascular pathway Netter, Frank H. "Mandibular Nerve." Netter Images, 2026, www.netterimages.com/mandibular-nerve-labeled-cleland-1e-general-anatomy-frank-h-netter-10792.html.

B. Mental Foramen and Anterior Loop

[10 pt Regular, Fully Justified, Indented First Line]

The mental foramen is usually located near the premolar region, but its position and anterior loop vary between patients. Implants placed in the second premolar region may traumatize the mental nerve if the foramen is not clearly identified. Injury may lead to pain, sensory reduction, hematoma, or persistent altered sensation in the lower lip and chin. Cross-sectional imaging is particularly important when the planned implant is close to the premolar zone.

C. Lingual Cortical Plate and Floor-of-Mouth Vessels

Perforation of the lingual cortical plate can injure sublingual or submental vessels and may cause rapidly expanding hematoma in the floor of the mouth. This complication is clinically significant because swelling in the sublingual and submental spaces may elevate the tongue and compromise the airway. The risk is higher when the mandibular ridge has a lingual concavity or when osteotomy angulation is directed lingually.

D. Dense Cortical Bone and Thermal Osteonecrosis

Mandibular cortical bone is denser than maxillary bone and may generate more heat during drilling. Excessive thermal trauma can cause osteonecrosis, delayed osseointegration, pain, or implant failure. Prevention depends on sharp drills, staged drilling, controlled speed, intermittent pressure, and copious external or internal irrigation.




Figure 4. Mandibular neurovascular anatomy in relation to implant placement. Hansen, John T. "Head and Neck." Card 1.7 in Netter's Anatomy Flash Cards. 3rd ed. Philadelphia: Saunders Elsevier, 2010.

V. ARTERIAL ARCHITECTURE AND SURGICAL HEMORRHAGE

The external carotid artery supplies the maxillofacial region through branches including the facial, lingual, and maxillary arteries. The maxillary artery is particularly important because its mandibular, pterygoid, and pterygopalatine segments supply deep structures involved in implant surgery. The inferior alveolar artery accompanies the inferior alveolar nerve through the mandibular canal; the posterior superior alveolar arteries supply the posterior maxilla and sinus wall;

and the descending palatine and sphenopalatine branches contribute to palatal and nasal vascular networks.

Aortic arch and carotid branching patterns explain the major inflow to the head and neck. The ophthalmic circulation also communicates with terminal branches of the facial artery through the angular and dorsal nasal arteries. These anastomoses are relevant not because implant surgery routinely enters them, but because they illustrate the dense vascular continuity that characterizes the midface.




Figure 5. Aortic arch and carotid-subclavian branching pattern.
Neck - Atlas of Anatomy. (2026). Doctorlib.Org. <https://doctorlib.org/medical/anatomy/39.html>




Figure 6. External carotid arterial branches relevant to maxillofacial surgery Arteries to Brain and Meninges. (2026). Netterimages.Com

VI. NEURAL ANATOMY AND POSTOPERATIVE NEUROSENSORY ASSESSMENT

The trigeminal nerve provides the main sensory innervation of the maxillofacial region. The maxillary division contributes infraorbital and superior alveolar branches, while the mandibular division gives rise to the inferior alveolar, mental, lingual, buccal, and auriculotemporal nerves. Implant-related neural complications may be caused by

mechanical trauma, compression from hematoma, inflammatory edema, chemical irritation, or thermal injury. Postoperative sensory changes should be documented early using a structured neurosensory examination. Symptoms such as numbness, burning, electric-shock pain, altered lip sensation, or tongue sensory changes should prompt radiographic review of implant position and assessment for compression, canal violation, or infection.

TABLE I. CLINICAL RISK MATRIX FOR SURGICAL PLANNING

Anatomical Zone	Principal Complication	Preventive Measure
Posterior maxilla	Sinus membrane perforation, sinusitis, implant migration	CBCT analysis of sinus floor, membrane, septa, and residual bone height
Anterior maxilla	Nasopalatine canal injury, pain, sensory alteration	Identification of canal morphology and palatal implant positioning when indicated

Topographic and Neurovascular Determinants of Dental Implant Complications: An Anatomical Review for Surgical Planning

Premolar mandible	Mental nerve trauma, lower-lip paresthesia	CBCT localization of mental foramen and anterior loop
Posterior mandible	Inferior alveolar nerve injury	Maintenance of neurovascular safety margin and controlled osteotomy depth
Lingual mandible	Sublingual/submental hematoma, airway risk	Assessment of lingual concavity and avoidance of lingual perforation
Dense mandibular cortex	Thermal osteonecrosis	Sharp drills, controlled speed, minimal pressure, and continuous irrigation

VII. DISCUSSION

A risk-zone approach shifts implant planning from a purely prosthetic or bone-volume analysis toward a multidimensional surgical safety model. The maxilla is dominated by sinus, thin-cortex, esthetic, and vascular-anastomotic risks. The mandible is dominated by neurovascular, lingual-perforation, and thermal-injury risks. This distinction explains why a technically similar osteotomy can have different consequences in the two jaws. Recent clinical data highlight the significant prevalence of implant positioning errors and their association with these specific anatomical complications [6].

CBCT is central to this framework because it permits visualization of three-dimensional anatomy that cannot be reliably inferred from panoramic imaging alone. However, imaging is only protective when it is translated into operative decisions: implant length and diameter selection, guided or freehand trajectory, safety margins, flap design, irrigation protocol, and postoperative monitoring.

VIII. CONCLUSION

Maxillofacial anatomy is not a passive background for implant placement; it actively determines the pathogenesis of surgical, vascular, infectious, and neurogenic complications. A journal-ready anatomical review should therefore emphasize the maxilla and mandible as distinct risk environments. The most important preventive principles are CBCT-based planning, preservation of neurovascular and

sinus safety margins, careful control of osteotomy temperature, and early recognition of sensory or vascular warning signs.

REFERENCES

- I. Bali RK, Sharma P, Gaba S, Kaur A, Ghanghas P. A review of complications of odontogenic infections. *National Journal of Maxillofacial Surgery*. 2015;6(2):136-143.
- II. Scarano A, et al. Current status of peri-implant diseases: a clinical review for evidence-based practice. *Journal of Functional Biomaterials*. 2023;14(5):210.
- III. Neal TW, Schlieve T. Complications of severe odontogenic infections: a review. *Biology*. 2022;11(12):1784.
- IV. Peña-Cardelles JF, et al. Inferior alveolar nerve damage related to dental implant placement: systematic review and meta-analysis. *Journal of Oral Implantology*. 2025.
- V. Sghaireen MG, et al. A CBCT-based three-dimensional assessment of the posterior mandible for implant planning. *Diagnostics*. 2020;10(6):406.
- VI. Alaqeely R, et al. Prevalence of dental implant positioning errors and related anatomical complications. *Journal of Clinical Medicine*. 2025;14(9):3221.